



Figure 3: Mean caloric intake and distribution of macronutrients (as percentage of total kilocalories) of participants during year 1 and year 4 of medical school. * $P = 0.02$, change from year 1 to year 4; $†P < 0.01$, change from year 1 to year 4; $‡P < 0.001$, change from year 1 to year 4. Stripes represent carbohydrate; dots represent protein; solid black represents fat; solid white represents alcohol

participants with blood pressure levels indicating hypertension. Still, 18.5% (18/97) of the study population in year 4 exhibited prehypertension, which is having systolic blood pressure between 120 and 139 mmHg or diastolic blood pressure between 80 and 89 mmHg. Elevated fasting insulin indicates insulin resistance, a risk factor for diabetes and cardiovascular disease. Although insulin levels increased significantly during the study period, mean fasting insulin values in both year 1 and year 4 were within the reference range.^[28] However, in year 4, 10 participants had insulin levels exceeding 25 $\mu\text{U/mL}$. Mean levels of HDL-cholesterol among participants were higher than the adult average^[37] and increased over the course of medical school. Lifestyle modification associated with increased HDL-cholesterol includes frequent aerobic activities. Interestingly, participants reported an increase in their physical activity during medical school, which may have been influenced by the opening of a fitness center within the medical college mid-way through the study period.

In our study, participants reported lifestyle habits congruent with national recommendations. Prior research has indicated that medical students report higher physical activity levels than age-matched peers in the general population.^[9] However, a recent longitudinal study using objective measures of physical fitness among medical students at the Uniformed Services University showed that muscular endurance and aerobic capacity declined during medical school.^[38] Our results concurred with these studies, showing a trend toward declining physical fitness during medical school, despite a reported increase in physical activity.

While college students often adopt unhealthy eating habits related to factors such as stress and lack of time, a recent study of Malaysian medical students by Ganasegeran *et al.*^[39] suggested that most medical students consume relatively healthy diets that meet national dietary guidelines. Similarly, our findings showed that participants' intake of carbohydrate, protein, and fat was within the acceptable macronutrient

distribution ranges, intake of most micronutrients met DRI recommendations, and HEI scores were indicative of a relatively high-quality diet in both years 1 and 4.

Despite the overall quality of the participants' diets, a few concerns were noted. With the increasing awareness of the essential role of Vitamin D in preventing osteoporosis and the common occurrence of insufficient Vitamin D status in the US, it is not surprising that Vitamin D intake increased during the study, particularly among females. Yet, the mean Vitamin D intake was below the current recommendation of 600 IU/day for both males and females throughout medical school. Similarly, some students may have been motivated to restrict sodium intake due to an emphasis on lowering dietary sodium for the prevention and treatment of hypertension. However, the average sodium intake of all participants remained higher than the American Heart Association recommendation of 1500-2300 mg/day. The role of trans fat in the promotion of cardiovascular disease led the National Heart, Lung, and Blood Institute and the American Heart Association to recommend consuming as little trans fat as possible. Yet, intake of trans fat, found primarily in convenience and processed foods, increased among males during the study period.

In a longitudinal survey of US medical students by Frank *et al.*,^[10] almost all students (97%) categorized their health as good, very good, or excellent, with a median of 1 day of poor physical health in the past month. Women reported significantly more days of poor mental health at the end of medical school (i.e. an increase from 1 to 2 days per month), while men reported no days of poor mental health at the start and finish of medical school. Our study results support the prior study's notion that medical students rate their health highly with very little occurrence of poor physical health. However, in our study, both males and females reported a higher frequency of poor mental health (median of 3 days per month) throughout medical school, perhaps related to multiple stressors such as academic challenges, financial concerns, or life events.^[11,12]

The results of several research studies suggest that the integration of nutrition and physical activity topics throughout the curriculum's didactic and clinical components is needed to positively influence students' knowledge and behaviors.^[4,40,41] During the study, in years 1 and 2 of medical school, the College's Clinical Foundations courses included nutrition lectures that were delivered by physicians and dietitians. The lecturers addressed dietary guidelines for disease prevention and nutrition therapy for several medical conditions, such as cardiovascular disease, diabetes, obesity, and critical illness. During the 3rd year clerkships, teaching strategies included lectures, case studies, online learning modules, and clinical experiences that addressed clinical nutrition. In addition, optional nutrition-related learning experiences were available to students. Clinical Nutrition, a 2-week elective course, was offered to 4th year students. The Medical Student Scholars Program – Nutrition Pathway was a longitudinal program that

offered nutrition-related extracurricular educational, clinical and research experiences to selected members who were admitted as 1st year students.

Limitations

There are inherent limitations to our study. Since all participants were recruited from one medical school, generalizability of results is limited to similar institutions. Enrollment in the study was voluntary and limited; therefore, study participants may not represent our medical student population in terms of health status or lifestyle habits. While we assessed alcohol consumption as part of the overall diet analysis, we did not collect further data on factors such as alcohol consumption patterns, smoking, or other coping behaviors which may have provided more insight into the participants' health. The longitudinal design and objective measurement of clinical parameters by trained health professionals following specific protocols are strengths of the study. Controversy remains over whether medical school does or does not affect the health status of students. Therefore, future longitudinal studies using similar clinical measures are needed with larger samples of medical students from several universities.

Conclusions

To the best of our knowledge, this is the first longitudinal study that assessed anthropometric and metabolic measures and lifestyle behaviors of medical students. Overall, our findings suggest that students' health and habits do not decline over the course of medical school; yet some students exhibit at-risk clinical measures, as well as diet and activity behaviors that are not aligned with national recommendations.

Prior research has shown that the health practices of medical students and physicians have the potential to play a highly influential role in counseling patients about disease prevention. Those who engage in healthy lifestyles are more likely to serve as role models and to provide quality counseling on exercise, diet, smoking, and other practices.^[4,42] Integration of the behavioral and social components of health (e.g., diet and physical activity) throughout medical school curricula has been recommended for over a decade by the Institute of Medicine^[43] and others.^[7,8] At the time of the study, the College of Medicine's inclusion of nutrition-related lectures, case studies, online modules, and clinical experiences throughout the curriculum and the addition of a fitness center are noteworthy. Such curricular initiatives that address essential lifestyle practices may be the first step in enhancing students' self-awareness, self-efficacy, and motivation to model healthy behaviors for their patients.^[40,42] Moreover, integration of behavioral counseling into the curriculum's didactic and experiential components may advance students' skills to engage in preventive counseling. Ultimately, such educational efforts may lead to healthier lifestyle habits and positive health outcomes for future physicians and their patients.

Acknowledgments

Amy Shova and Nicole Wrassmann, research assistants, provided valuable assistance with study coordination and data entry. The content of this paper is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

Financial support and sponsorship

The project was supported by the National Center for Advancing Translational Sciences of the National Institutes of Health, under Award Number UL1TR000077. Approval for the study was granted by the Institutional Review Boards of the University of Cincinnati (IRB 06-07-12-01) and Cincinnati Children's Hospital Medical Center (IRB 2008-1235).

Conflicts of interest

The salaries of Ms. Summer, Dr. Khoury, and Dr. Heubi are funded either entirely or in part by the National Center for Advancing Translational Sciences of the National Institutes of Health, under Award Number UL1TR000077, the grant which supported this study. Dr. Brehm, Dr. Filak, and Dr. Lieberman have no conflicts of interest related to this study.

References

1. Institute of Medicine. For the Public's Health: Investing in a Healthier Future. Washington, DC: The National Academies Press; 2012.
2. Frank E, Segura C, Shen H, Oberg E. Predictors of Canadian physicians' prevention counseling practices. *Can J Public Health* 2010;101:390-5.
3. Howe M, Leidel A, Krishnan SM, Weber A, Rubenfire M, Jackson EA. Patient-related diet and exercise counseling: Do providers' own lifestyle habits matter? *Prev Cardiol* 2010;13:180-5.
4. Lobelo F, Duperly J, Frank E. Physical activity habits of doctors and medical students influence their counselling practices. *Br J Sports Med* 2009;43:89-92.
5. Eisenberg DM, Burgess JD. Nutrition education in an era of global obesity and diabetes: Thinking outside the box. *Acad Med* 2015;90:854-60.
6. Phillips E, Pojednic R, Polak R, Bush J, Trilk J. Including lifestyle medicine in undergraduate medical curricula. *Med Educ Online* 2015;20:26150.
7. Kushner RF, Van Horn L, Rock CL, Edwards MS, Bales CW, Kohlmeier M, *et al.* Nutrition education in medical school: a time of opportunity. *Am J Clin Nutr* 2014;99 5 Suppl: 1167S-73S.
8. Torabi MR, Tao R, Jay SJ, Olcott C. A cross-sectional survey on the inclusion of tobacco prevention/cessation, nutrition/diet, and exercise physiology/fitness education in medical school curricula. *J Natl Med Assoc* 2011;103:400-6.
9. Frank E, Tong E, Lobelo F, Carrera J, Duperly J. Physical activity levels and counseling practices of U.S. medical students. *Med Sci Sports Exerc* 2008;40:413-21.
10. Frank E, Carrera JS, Elon L, Hertzberg VS. Basic demographics, health practices, and health status of U.S. medical students. *Am J Prev Med* 2006;31:499-505.

11. MacLean L, Booza J, Balon R. The impact of medical school on student mental health. *Acad Psychiatry* 2016;40:89-91.
12. Lins L, Carvalho FM, Menezes MS, Porto-Silva L, Damasceno H. Health-related quality of life of students from a private medical school in Brazil. *Int J Med Educ* 2015;6:149-54.
13. Majra J. Do our medical colleges inculcate health-promoting lifestyle among medical students: A pilot study from two medical colleges from Southern India. *Int J Prev Med* 2013;4:425-9.
14. Kjeldstadli K, Tyssen R, Finset A, Hem E, Gude T, Gronvold NT, *et al.* Life satisfaction and resilience in medical school – A six-year longitudinal, nationwide and comparative study. *BMC Med Educ* 2006;6:48.
15. Frank E, Carrera JS, Elon L, Hertzberg VS. Predictors of US medical students' prevention counseling practices. *Prev Med* 2007;44:76-81.
16. Brehm BJ, Lattin BL, Summer SS, Boback JA, Gilchrist GM, Jandacek RJ, *et al.* One-year comparison of a high-monounsaturated fat diet with a high-carbohydrate diet in type 2 diabetes. *Diabetes Care* 2009;32:215-20.
17. Baecke JA, Burema J, Frijters JE. A short questionnaire for the measurement of habitual physical activity in epidemiological studies. *Am J Clin Nutr* 1982;36:936-42.
18. Pols MA, Peeters PH, Bueno-De-Mesquita HB, Ocké MC, Wentink CA, Kemper HC, *et al.* Validity and repeatability of a modified Baecke questionnaire on physical activity. *Int J Epidemiol* 1995;24:381-8.
19. Ryhming I. A modified Harvard step test for the evaluation of physical fitness. *Arbeitsphysiologie* 1953;15:235-50.
20. Sharkey BJ. *Physiology of Fitness*. Champaign, IL: Human Kinetics; 1984.
21. DeVries HA. *Laboratory Experiments in Exercise Physiology*. Dubuque, Iowa: Wm. C. Brown; 1971.
22. Willett W. *Nutritional Epidemiology*. 3rd ed. Oxford: Oxford University Press; 2013.
23. Landy DC, Kurtz JM, Miller TL, Ludwig DA. Statistical program to automate the creation of Healthy Eating Index scores using Nutrition Data System for Research output. *J Acad Nutr Diet* 2012;112:A14.
24. Newschaffer C. Validation of Behavioral Risk Factor Surveillance System (BRFSS) HRQOL Measures in a Statewide Sample. Atlanta, Georgia: U.S. Department of Health and Human Services, Public Health Service, Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion; 1998.
25. Andresen EM, Catlin TK, Wyrwich KW, Jackson-Thompson J. Retest reliability of surveillance questions on health related quality of life. *J Epidemiol Community Health* 2003;57:339-43.
26. Jensen MD, Ryan DH, Donato KA, Apovian CM, Ard JD, Comuzzie AG, *et al.* Executive Summary: Guidelines (2013) for the management of overweight and obesity in adults. *Obesity* 2014 Jul; 22 Suppl 2:S5-39.
27. James PA, Oparil S, Carter BL, Cushman WC, Dennison-Himmelfarb C, Handler J, *et al.* 2014 evidence-based guideline for the management of high blood pressure in adults: Report from the panel members appointed to the Eighth Joint National Committee (JNC 8). *JAMA* 2014;311:507-20.
28. Melmed S, Polonsky KS, Larsen PR, Kronenberg HM. *Williams Textbook of Endocrinology*. Vol. 12. Philadelphia: Elsevier Saunders; 2011.
29. Heo M, Faith MS, Pietrobelli A, Heymsfield SB. Percentage of body fat cutoffs by sex, age, and race-ethnicity in the US adult population from NHANES 1999-2004. *Am J Clin Nutr* 2012;95:594-602.
30. Stone NJ, Robinson JG, Lichtenstein AH, Bairey Merz CN, Blum CB, Eckel RH, *et al.* 2013 ACC/AHA guideline on the treatment of blood cholesterol to reduce atherosclerotic cardiovascular risk in adults: A report of the American College of Cardiology/American Heart Association task force on practice guidelines. *J Am Coll Cardiol* 2014;63:2889-934.
31. Mustelin L, Joutsi J, Latvala A, Pietiläinen KH, Rissanen A, Kaprio J. Genetic influences on physical activity in young adults: A twin study. *Med Sci Sports Exerc* 2012;44:1293-301.
32. Camhi SM, Katzmarzyk PT. Total and femoral neck bone mineral density and physical activity in a sample of men and women. *Appl Physiol Nutr Metab* 2012;37:947-54.
33. National Institutes of Health, Office of Dietary Supplements. Nutrient Recommendations: Dietary Reference Intakes. Available from: https://www.ods.od.nih.gov/Health_Information/Dietary_Reference_Intakes.aspx. [Last accessed on 2016 May 01].
34. Ervin RB. Healthy Eating Index-2005 Total and Component Scores for Adults Aged 20 and Over: National Health and Nutrition Examination Survey, 2003-2004. National Health Statistics Reports. Vol. 44; 2011. p. 1-9.
35. Ogden CL, Carroll MD, Kit BK, Flegal KM. Prevalence of childhood and adult obesity in the United States, 2011-2012. *JAMA* 2014;311:806-14.
36. Nwankwo T, Yoon SS, Burt V, Gu Q. Hypertension among adults in the United States: National Health and Nutrition Examination Survey, 2011-2012. NCHS Data Brief 2013;133:1-8.
37. Carroll MD, Kit BK, Lacher DA, Shero ST, Mussolino ME. Trends in lipids and lipoproteins in US adults, 1988-2010. *JAMA* 2012;308:1545-54.
38. Stephens MB, Cochran C, Hall JM, Olsen C. Physical fitness during medical school: A 4-year study at the Uniformed Services University. *Fam Med* 2012;44:694-7.
39. Ganasegeran K, Al-Dubai SA, Qureshi AM, Al-Abed AA, Am R, Aljunid SM. Social and psychological factors affecting eating habits among university students in a Malaysian medical school: A cross-sectional study. *Nutr J* 2012;11:48.
40. Dacey ML, Kennedy MA, Polak R, Phillips EM. Physical activity counseling in medical school education: A systematic review. *Med Educ Online* 2014;19:24325.
41. Kris-Etherton PM, Akabas SR, Bales CW, Bistrrian B, Braun L, Edwards MS, *et al.* The need to advance nutrition education in the training of health care professionals and recommended research to evaluate implementation and effectiveness. *Am J Clin Nutr* 2014;99 5 Suppl: 1153S-66S.
42. Frank E, Elon L, Hertzberg V. A Quantitative assessment of a 4-year intervention that improved patient counseling through improving medical student health. *MedGenMed* 2007;9:58.
43. Institute of Medicine. *Improving Medical Education: Enhancing the Behavioral and Social Science Content of Medical School Curricula*. Washington, DC: The National Academies Press; 2004.