

Clinical and Radiographic Evaluation of Stainless Steel Crowns and Zirconia Crowns in Primary Molars: A Systematic Review

Deepa Gurunathan* and Sruthi Suguna

Department of Pediatric and Preventive dentistry, Saveetha Dental College and hospitals, Saveetha Institute of Medical and Technical sciences, Chennai, India

Corresponding author:

Dr. Deepa Gurunathan. Department of Pediatric and Preventive dentistry, Saveetha Dental college and hospitals, Saveetha University, Chennai, India, Tel:9994619386; E-mail: deepag@saveetha.com

Abstract

Aims: To systematically identify and evaluate the clinical and radiographic success of stainless steel crown and zirconia crown in primary mandibular molars. **Materials & Methods:** A comprehensive search was done using electronic databases such as PubMed Central, Cochrane Database of Systematic Reviews, Lilacs, Science Direct and SIGLE. The title scan was done to identify relevant articles, which were further evaluated for inclusion by reading the abstract. Bibliography of all the identified studies was scanned to include studies published outside the electronic database. All the studies comparing the clinical and radiographic evaluation of stainless steel crowns and zirconia crowns in primary molars were included. **Results:** Electronic database search identified 205 articles. Out of which, 187 were excluded after reading titles and removing duplication. The 8 studies were evaluated in detail after reading the abstract and full text. A final of 2 studies were included based on the inclusion criteria to meet the research question. Both the studies analysed only the clinical parameters of stainless steel crowns and zirconia crowns in primary molars whereas radiographic evaluation was not mentioned. **Conclusion:** Limited studies are available comparing the clinical and radiographic success of stainless steel crowns and zirconia crowns in primary mandibular molars. The current systematic review identifies the need for more high quality clinical trials evaluating the clinical and radiographic parameters of stainless steel crowns and zirconia crowns in primary mandibular molars.

Keywords: Stainless steel; Crowns; Zirconia; Mandibular molars; systematic review

Introduction

Oral health plays a fundamental role in total health and well-being. It influences numerous aspects of a person's health status and quality of life including self-esteem, learning, employment, ability to masticate, speak and other levels of routine activities. Early Childhood Caries (ECC) has been studied to be the most prevalent chronic disease of babyhood. [1] ECC is defined as "the existence of one or more decayed, missing (due to caries), or filled tooth surfaces in any primary tooth in a child 71 months of age or younger." [2] As ECC is of aggressive nature, cavitation may develop rapidly due to the areas of demineralization and hypoplasia. If the disease is left untreated, it may involve the pulp, which often leads to dental infection and possibly life-threatening fascial space involvement. Such dental infections may require emergency medical care such as hospitalization, antibiotics and probably extraction of the affected tooth. [3] Children those are at high risk for ECC should have a caretaker appointed by a practitioner who has skills, training, experience, and expertise to manage both the clinical and the disease process. To execute the treatment safely, effectively, and efficiently, the pediatric dentist should administer advanced behavior guidance techniques, protective stabilization and/or sedation or general anesthesia. [4]

Stainless Steel Crowns (SSCs) has been employed for the restoration of primary teeth that are damaged/affected by caries,

decalcification around the neck of the tooth, and developmental anomalies (e.g., hypoplasia, hypo calcification). It is also used when the downfall of further accessible restorative supplies is more probable (e.g., interproximal caries ranging farther than line angles, children with bruxism). Moreover, next to pulpotomy or pulpectomy, SSC may be used for restoration of a primary tooth which will be exploited as an abutment to maintain space or to be used as interposed rehabilitation of severed teeth. [5] Usage of SSCs should also be considered in patients with increased risk of caries whose participation is distressed by age, behavior, or medical history. Most often, these patients undergo treatment under sedatives or general anesthetics. [6-12] SSCs tend to last longer in patients with developmental or medical conditions that do not improve as they age, thus reducing the possible usage of sedatives and general anesthetics considering that it is costly and its inherent perils. However then again, severely damaged primary dentitions among pediatric patients is deemed to be one with immense difficulties, to treat esthetically. [13-19] During the past 50 years, the prominence of taking care of mostly degenerating primary dentition changed from removal to

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 3.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as the author is credited and the new creations are licensed under the identical terms.

How to Cite this Article: Gurunathan D, et al. Clinical and Radiographic Evaluation of Stainless Steel Crowns and Zirconia Crowns in Primary Molars: A Systematic Review. Ann Med Health Sci Res. 2021;11:11-17.

rehabilitation. Early recovery includes employment of stainless steel binding or crowns on critically damaged dentition. While working, they were unaesthetic, and its usage is confined to posterior teeth. The mesio-buccal area of the first primary molars and second maxillary primary molars may be seen when the child smiles.

During the past 20 years, there was an increased demand by adults in the esthetic restoration of their jeopardized dentures. Equivalently, a higher esthetic standard is expected by parents for the recovery of their children's carious teeth. More recently, Zirconia esthetic crowns appeared on the market.^[20-22] Zirconia is a crystalline dioxide of zirconium that has mechanical properties akin to those of metals, and its color is similar to that of teeth. Ready-made zirconia crowns are available for both, primary incisors and molars. Our department is passionate about child care, we have published numerous high quality articles in this domain over the past 3 years previously our team has a rich experience in working on various research projects across multiple disciplines.^[24-38] Now the growing trend in this area motivated us to pursue this project.

With this inspiration we planned to pursue a systematic review on clinical and radiographic success of SSC and

Methodology

This review was done in accordance with guidelines given by Cochrane Handbook of Systematic Review.

Search strategy

The controlled vocabulary (MeSH terms) and free words in the search strategy were defined based on the Accepted Article following PICOS strategy.

Population: Primary molars

Intervention: Zirconia crowns

Comparison: Stainless steel crowns

Outcome: Clinical and Radiographic success

The search strategy was initially established for the PubMed database, associating controlled vocabulary (MeSH terms) and free-words. The boolean operator OR was used to combine the terms in each PICO concept; the operator AND were used to combine the different PICO concepts (population, intervention, and comparison). The strategy was adapted to other electronic databases (Scopus, Web of Science, Latin American Literature of Health Sciences of the Americas and Caribbean - LILACS, Brazilian Library of Dentistry). The grey literature was searched using the databases System for Information on Grey Literature in Europe (SIGLE) and Scholar Google. (Primary molar or Deciduous teeth) and (Zirconia crown or Tooth coloured crown) and (Stainless steel crown or Metal crown or Pre-formed metal crown) and (Clinical success or Radiographic success).

Hand searches were undertaken to find additional relevant published material that might have been missed in electronic searches. The articles published till October 2019 were included in the present study.

Eligibility criteria

Eligible study designs in paediatric patients up to 10 years of age receiving crown placement, randomized controlled studies comparing the clinical and radiographic success of both the crowns and studies published in English language only were included in the study. Studies involving older children or adults, studies involving comparison with pre-veneered stainless steel crowns, ongoing studies in which results have not yet been published, case series, descriptive and *in vitro* studies were not included.

Data collection and analysis

One author carried out the search strategy for the individual databases. The total number of titles obtained were scanned and evaluated independently by two authors. The studies duplicated in the different databases were excluded. In case of any disagreement between the two authors, a final decision was obtained by discussion between the two authors. Abstracts of the studies were evaluated when complete information regarding the groups and participants included was not mentioned in the title. The abstract evaluation was carried out independently by two authors to identify the final studies to be included based on the inclusion and exclusion criteria. Full text articles were evaluated when the abstracts did not provide adequate information regarding the groups compared. Figure 1 gives the PRISMA flow diagram for the included studies [Figure 1]. The final studies included by the discussion of both the authors were further evaluated for the quality of studies following the guidelines given by Cochrane Handbook of systematic review. This was done independently by both the authors and any discrepancy was resolved by discussion between both the authors.

The Cochrane Collaboration tool for assessing risk of bias in randomized trials was used for the quality assessments of the trials, following the recommendations described in the Cochrane Handbook for Systematic Reviews of Interventions 5.1.0.^[39,40] This was accomplished by two independent reviewers. The Cochrane tool is based on six domains: adequate sequence generation, allocation concealment, blinding of the outcome assessors, incomplete outcome data, selective outcome reporting, and other possible sources of bias. The judgment for each domain consisted of recording "yes" (low risk of bias), "no" (high risk of bias) or "unclear" (either lack of information or uncertainty about the potential for bias). Two out of the six domains in the Cochrane risk of bias tool were considered as key domains for this systematic review (sequence generation and allocation concealment).^[39] The papers were judged to be at "low" risk of bias if they were judged as "low" risk in both key domains. If one key domain was classified as "unclear" or "high" risk of bias, the study was considered at "unclear Accepted Article" or "high" risk of bias, respectively.

Study Selection

The systematic search from the electronic databases of PubMed revealed 23 studies, Cochrane library revealed 33 studies, Google Scholar revealed 1 study and Science Direct revealed 147




Figure 1: Gives the prisma flow diagram for the included studies. The final studies included by the discussion of both the authors were further evaluated for the quality of studies following the guidelines given by Cochrane handbook of systematic review. This was done independently by both the authors and any discrepancy was resolved by discussion between both the authors.

studies. No studies were obtained from the database of LILACS and SIGLE and 1 study was obtained from hand searching. After removal of duplicates and title scan, 8 studies were identified. After abstract scanning 6 articles were eliminated as they did not meet the inclusion and exclusion criteria. Full text articles for the other 2 studies were obtained for more detailed evaluation. The bibliography of these full text articles was scanned to include studies apart from the electronic databases. No relevant studies were found from the cross-reference. A total of 2 studies met the inclusion and exclusion criteria of the intended research.

Study Characteristics

The following studies by Kevin et al. and kinay et al. were included due to similarities in the study design, intervention and control groups and the parameters observed and were evaluated.^[41,42] The studies by Kratunoya et al.; Walia; Leith et al.; Yilmaz et al.; kara et al. were excluded because they did not meet the inclusion criteria as studies were made up of comparison of SSCs with crowns other than zirconia.^[43-47] The risk of bias in the studies is shown [Figures 2 and 3] Kevin et al., assessed the clinical and radiographic success of children




Figure 2: Risk of bias summary: Judgement about each risk of bias item for each included study.




Figure 3: Risk of bias Graph: Judgments about each risk of bias item presented as percentage among included studies.

steel and zirconia crowns and found no clinically significant differences between the two crowns in relation to gingival health, restoration success, opposing tooth wear, anatomic form, marginal integrity, marginal discoloration, proximal contact area and secondary caries but there was a significant difference found related to parental satisfaction with zirconia crowns.^[41]

Taran et al. assessed the clinical and radiographic success of children receiving stainless steel and zirconia crowns and found that teeth restored with zirconia crowns showed less plaque accumulation and better gingival health as compared to teeth with stainless steel crowns. She also observed that both stainless steel crowns and zirconia crowns used in primary dentition had successful clinical outcomes while a minority of zirconia crowns show minor staining and fracture.^[42]

Discussion

The preservation of healthy primary dentition is important for the overall development of children. Treating children affected by severe ECC is extremely difficult for the pediatric dentist. The preservation of tooth structure, strength and longevity of the restoration, and parental satisfaction are three aspects to be considered. In recent years, technological advancements in dental materials for children have necessitated reassessments of new treatment options. Various approaches have been emerged for full-coverage restorations in recent times.^[48] Since 1950, Stainless Steel Crowns (SSCs) have been the most frequently preferred full-coverage restoration for the primary and permanent posterior teeth of children.^[49] The outgrowth of SSCs in preserving severely decayed or pulp-tomized primary molars have been reported in the literature.^[18] Although SSCs are low maintenance, their metallic appearance has been a controversial issue for the esthetical perceptions of parents.^[50] Today, esthetic restoration of primary teeth is becoming as important as their durability in pediatric dentistry. The metallic appearance of SSCs is frequently raised as an issue by children and parents.^[51,52] The recently developed prefabricated Zirconia Crowns (ZCs) are a fairly new treatment option that enables practitioners to provide superior esthetic results. Although zirconia has been accepted as a restorative material in permanent dentition for many years, its use in the primary dentition started in 2008.^[53,54] Zirconia crowns have mechanical properties similar to those of metal but do not allow crimping or contouring and require extensive tooth reduction.^[55]

Clinical success has continued to be a point of discussion with the development of new restorative options. There were only a few randomized clinical trials to compare Zirconia crowns

receiving stainless with SSCs that are considered to be the treatment of reference, especially on primary molars. Hence the aim of this study was to investigate the outcome of both the crowns using the available data.

According to the current systematic review, study done by Kevin et al., showed that gingival health appeared to be better adjacent to zirconia crowns at 6 months, compared to SSCs; however, at 2 years, the gingiva appeared similar adjacent to either crown.^[48]

Kinay et al. found the molars treated with zirconia crowns showed better periodontal index and gingival index scores than those treated with SSCs.^[49] Abdulhadi et al., showed that gingival health was better in teeth restored with zirconia crowns than in those which treated with SSCs during 3 and 6 months follow up, but 12 months follow-up revealed that both groups presented with healthy gingiva. These results could be explained as zirconia used for tooth component exhibits remarkable biocompatibility as well as shows smooth plus polished exterior which lead to lower tendency of plaque build-up and thus lower chance of gingival irritation.^[56] Sharaf et al. revealed that SSCs crowns had poorly adapted margins showing signs of gingivitis while variations in crown marginal extension and adequacy of the crowns as judged radiographically had no effect on the gingival health.^[57] This agrees with the work of Henderson, Myers and Checcio et al who reported high incidence of gingivitis around poorly fitting and improperly contoured crowns due to failure of maintaining a clean area around such crowns.^[58-61] It was the opinion of Myers et al. that SSCs should be carefully fitted to avoid mechanical irritation to the gingival tissues.^[61] On the other hand, Webber found no relation better SSCs and gingival health.^[62] Kara et al. observed a decrease in oral hygiene levels in their study and suggested that oral hygiene instructions be given to children whose teeth were restored with prefabricated crown.^[47] According to our review, Kinay et al. suggested that the children should receive standard oral hygiene education as a part of their routine care. This instruction along with frequent follow-up appointments might have motivated the children to maintain a good level of oral hygiene throughout their study.

A study done by Sharaf et al. revealed that oral hygiene factor affected gingival health. Children with poor oral hygiene showed higher incidence of gingivitis while children with good oral hygiene maintained healthy gingiva around SSCs.^[26] These findings are in agreement with Checcio et al who concluded that individuals with poor oral hygiene showed pronounced tissue degeneration despite the quality of SSCs and that improperly contoured restorations predispose the gingiva to more severe inflammation.^[60]

Kevin et al postulated that the wear of the opposing arch tooth appeared higher in zirconia group at 6 months; by two years, however, there was no difference between the zirconia and SSCs in opposing tooth wear.^[48] A slight tooth wear in natural dentition is considered normal.^[63] If restorative dental materials have different wear properties compared to the natural teeth, they can affect the wear rate of antagonist natural teeth.^[64] Therefore, wear occurring between enamel and dentin restorations is an important factor that should always be considered in the selection of restorative materials in clinical practice.

Alveolar bone loss was associated with SSCs that were judged

radiographically as non-satisfactory^[55] which was in agreement with the literature as Beimstein et al. who reported an association between alveolar bone resorption and inadequately placed SSCs and also in sites adjacent to proximal caries.^[65] Again, Guelman et al. stated that a well-adapted crown in primary second molar does not facilitate the appearance of marginal alveolar bone loss on the adjacent first permanent molar.^[66] Our institution is passionate about high quality evidence based research and has excelled in various fields. We hope this study adds to this rich legacy.

Research journals on outcome measurements are very crucial for clinical day-to-day processes as they may administer clinicians with better understanding for making appropriate treatment opinions. They also furnish appropriate and more accurate information in order to gain valid consent from the patients. The consolidation of the outcomes of multiple research papers, such as in the form of meta-analyses, can produce very powerful conclusions. The results of this systematic review showed similar outcomes in both the studies.^[67-72]

Conclusion

Zirconia crowns had a considerable advantage over SSCs due to their esthetic appearance. Teeth restored with zirconia crowns showed less plaque accumulation and better gingival health than teeth restored with SSCs. Both SSCs and zirconia crowns can be used in primary dentition with successful clinical outcomes, while a minority of zirconia crowns show minor staining and fracture. Further randomized controlled trials, with greater numbers conducted over a longer period of time, would be of benefit in improving our understanding of this subject. Based on the current data, clinicians need to be more cautious about using the crowns to restore posterior teeth in primary molars.

Conflict of Interest

The authors and planners have disclosed no potential conflicts of interest, financial or otherwise.

Acknowledgements

The authors would like to acknowledge all volunteers who participated in the study.

References

- Retnakumari N, Cyriac G. Childhood caries as influenced by maternal and child characteristics in pre school children of Kerala-an epidemiological study. *Contemp Clin Dent*. 2012;3:2–8.
- American academy of pediatrics. Policy on early childhood caries (ECC): Classifications, consequences, and preventive strategies. *Pediatr Dent*. 2008;30:40–3.
- Sheller B, Williams BJ, Lombardi SM. Diagnosis and treatment of dental caries-related emergencies in a children's hospital. *Pediatr Dent*. 1997;19:470–5.
- Pediatric restorative dentistry. *Pediatr Dent*. 2017;39(6):312–24.
- Ravikumar DNS, Ramakrishna M, Sharna N, Robindro W. Evaluation of McNamara's analysis in South Indian (Tamil Nadu) children between 8-12 years of age using lateral cephalograms. *J Oral Biol Craniofac Res*. 2019;9:193–7.
- Ravikumar D, Gurunathan D, Gayathri R, Priya V, Geetha RV. DNA profiling of *Streptococcus mutans* in children with and without black tooth stains: A polymerase chain reaction analysis. *Dent Res J*. 2018;15:334.
- Ravikumar D, Jeevanandan G, Subramanian EMG. Evaluation of knowledge among general dentists in treatment of traumatic injuries in primary teeth: A cross-sectional questionnaire study. *Eur J Dent*. 2017;11:232–7.
- Ravindra V, Rekha V, Annamalai S, Sharmin D, Norouzi-Baghkomeh P. A comparative evaluation between dermatoglyphic patterns and different terminal planes in primary dentition. *J Clin Exp Dent*. 2018;10:e1149–54.
- Ravindra V, Rekha CV, Annamalai S, Sharmin DD, Norouzi-Baghkomeh P. A comparative evaluation between cheiloscopy patterns and the permanent molar relationships to predict the future malocclusions. *J Clin Exp Dent*. 2019;11:e553–7.
- Govindaraju L, Jeevanandan G, Subramanian EMG. Comparison of quality of obturation and instrumentation time using hand files and two rotary file systems in primary molars: A single-blinded randomized controlled trial. *Eur J Dent*. 2017;11:376–9.
- Govindaraju L, Jeevanandan G, Subramanian EMG. Knowledge and practice of rotary instrumentation in primary teeth among indian dentists: A questionnaire survey. *J Int Oral Health*. 2017;9:45.
- Nair M, Jeevanandan G, Vignesh R, Emg S. Comparative evaluation of post-operative pain after pulpectomy with k-files, kedo-s files and mtwo files in deciduous molars -A randomized clinical trial. *BDS*. 2018 24;21:411.
- Jeevanandan G, Ganesh S, Arthilakshmi. Kedo file system for root canal preparation in primary teeth. *Indian J Dent Res*. 2019;30:622–4.
- Panchal V, Jeevanandan G, Subramanian E. Comparison of instrumentation time and obturation quality between hand K-file, H-files, and rotary Kedo-S in root canal treatment of primary teeth: A randomized controlled trial. *J Indian Soc Pedod Prev Dent*. 2019;37:75–9.
- Subramanyam D, Gurunathan D, Gaayathri R, Vishnu Priya V. Comparative evaluation of salivary malondialdehyde levels as a marker of lipid peroxidation in early childhood caries. *Eur J Dent*. 2018;12:67–70.
- Vignesh R, Sharmin D, Rekha CV, Annamalai S, Baghkomeh PN. Management of complicated crown-root fracture by extra-oral fragment reattachment and intentional reimplantation with 2 years review. *Contemp Clin Dent*. 2019;10:397–401.
- Ramadurai N, Gurunathan D, Samuel AV, Subramanian E, Rodrigues SJL. Effectiveness of 2% Articaine as an anesthetic agent in children: Randomized controlled trial. *Clin Oral Investig*. 2019;23:3543–50.
- Panchal V, Gurunathan D, Shanmugaavel AK. Smartphone application as an aid in determination of caries risk and prevention: A pilot study. *Eur J Dent*. 2017;11:469–74.
- Panchen V, Jeevanandan G, Subramanian EMG. Comparison of post-operative pain after root canal instrumentation with hand K-files, H-files and rotary Kedo-S files in primary teeth: A randomized clinical trial. *Eur Arch Paediatr Dent*. 2019;20:467–72.
- Jeevanandan G, Govindaraju L. Clinical comparison of Kedo-S pediatric rotary files vs. manual instrumentation for root canal preparation in primary molars: A double blinded randomized clinical trial. *Eur Arch Paediatr Dent*. 2018;19:273–8.

21. Samuel SR, Acharya S, Rao JC. School Interventions-based Prevention of Early-Childhood Caries among 3-5-year-old children from very low socioeconomic status: Two-year randomized trial. *J Public Health Dent.* 2020;80:51–60.
22. Ramakrishnan M, Dhanalakshmi R, Subramanian EMG. Survival rate of different fixed posterior space maintainers used in Paediatric Dentistry - A systematic review. *Saudi Dent J.* 2019;31:165–72.
23. Vishnu Prasad S, Kumar M, Ramakrishnan M, Ravikumar D. Report on oral health status and treatment needs of 5-15 years old children with sensory deficits in Chennai, India. *Spec Care Dentist.* 2018;38:58–9.
24. Ponnulakshmi R, Shyamaladevi B, Vijayalakshmi P, Selvaraj J. In silico and *in vivo* analysis to identify the antidiabetic activity of beta sitosterol in adipose tissue of high fat diet and sucrose induced type-2 diabetic experimental rats. *Toxicol Mech Methods.* 2019;29:276–90.
25. Mathew MG, Samuel SR, Sonny AJ, Roopa KB. Evaluation of adhesion of *Streptococcus mutans*, plaque accumulation on zirconia and stainless steel crowns, and surrounding gingival inflammation in primary molars: randomized controlled trial. *Clin Oral Investig.* 2020;24:3275–80.
26. Subramaniam N, Muthukrishnan A. Oral mucositis and microbial colonization in oral cancer patients undergoing radiotherapy and chemotherapy: A prospective analysis in a tertiary care dental hospital. *J Investig Clin Dent.* 2019;10:e12454.
27. Girija ASS, Shankar EM, Larsson M. Could SARS-CoV-2-Induced hyperinflammation magnify the severity of Coronavirus disease (Covid-19) leading to acute respiratory distress syndrome? *Front Immunol.* 2020;27:1206.
28. Dinesh S, Kumaran P, Mohanamurugan S, Vijay R, Singaravelu DL, Vinod A, et al. Influence of wood dust fillers on the mechanical, thermal, water absorption and biodegradation characteristics of jute fiber epoxy composites. *J Polym Res.* 2020;27.
29. Thanikodi S, Singaravelu D Kumar, Devarajan C, Venkatraman V, Rathinavelu V. Teaching learning optimization and neural network for the effective prediction of heat transfer rates in tube heat exchangers. *Therm Sci.* 2020;24:575–81.
30. Murugan MA, Jayaseelan V, Jayabalakrishnan D, Maridurai T, Kumar SS, Ramesh G, et al. Low velocity impact and mechanical behaviour of shot blasted SiC wire-mesh and silane-treated aloevera/hemp/flax-reinforced SiC whisker modified epoxy resin composites. *Silicon Chem.* 2020;12:1847–56.
31. Vadivel JK, Govindarajan M, Somasundaram E, Muthukrishnan A. Mast cell expression in oral lichen planus: A systematic review. *J Investig Clin Dent.* 2019;10:e12457.
32. Chen F, Tang Y, Sun Y, Veeraraghavan VP, Mohan SK, Cui C. 6-shogaol, a active constituents of ginger prevents UVB radiation mediated inflammation and oxidative stress through modulating Nrf2 signaling in human epidermal keratinocytes (HaCaT cells). *J Photochem Photobiol B.* 2019;197:111518.
33. Manickam A, Devarasan E, Manogaran G, Priyan MK, Varatharajan R, Hsu C-H, et al. Score level based latent fingerprint enhancement and matching using SIFT feature. *Multimed Tools Appl.* 2019;78:3065–85.
34. Wu F, Zhu J, Li G, Wang J, Veeraraghavan VP, Krishna Mohan S, et al. Biologically synthesized green gold nanoparticles from induce growth-inhibitory effect on melanoma cells (B16). *Artif Cells Nanomed Biotechnol.* 2019;47:3297–305.
35. Ma Y, Karunakaran T, Veeraraghavan VP, Mohan SK, Li S. Sesame inhibits cell proliferation and induces apoptosis through inhibition of STAT-3 translocation in thyroid cancer cell lines (FTC-133). *Biotechnol Bioprocess Eng.* 2019;24:646–52.
36. Ponnankajamdeen M, Rajeshkumar S, Vanaja M, Annadurai G. *In vivo* type 2 diabetes and wound-healing effects of antioxidant gold nanoparticles synthesized using the insulin plant *Chamaecostus cuspidatus* in albino rats. *Can J Diabetes.* 2019;43:82–9.e6.
37. Vairavel M, Devaraj E, Shanmugam R. An eco-friendly synthesis of *Enterococcus* sp.-mediated gold nanoparticle induces cytotoxicity in human colorectal cancer cells. *Environ Sci Pollut Res Int.* 2020; 27:8166–75.
38. Paramasivam A, Priyadharsini VJ, kumar RS. N6-adenosine methylation (m6A): a promising new molecular target in hypertension and cardiovascular diseases. *Hypertens Res.* 2020;43:153–4.
39. Higgins JPT, Altman DG, Gotzsche PC, Juni P, Moher D, Oxman AD, et al. The cochrane collaboration's tool for assessing risk of bias in randomised trials. *BMJ.* 2020;43:153–4.
40. Higgins JPT, Green S. *Cochrane handbook for systematic reviews of interventions.* Wiley. 2008;p:672.
41. Donly KJ, Sasa I, Contreras CI, Mendez MJC. Prospective randomized clinical trial of primary molar crowns: 24-month results. *Pediatr Dent.* 2018;40:253–8.
42. Kaya MS, Taran PK, Bakkal M. Clinical follow- up of occlusal vertical dimension increase in stainless steel crowns applied with hall technique: A pilot study. *Yeditepe Dental Journal.* 2018;37–42.
43. Kratunova E, O'Connell AC. Chairside repair of veneered primary molar stainless steel crowns: A pilot study. *Pediatr Dent.* 2015;37:46–50.
44. Salami A, Walia T, Bashiri R. Comparison of parental satisfaction with three tooth-colored full-coronal restorations in primary maxillary incisors. *Journal of Clinical Pediatric Dentistry.* 2015;pp:423–8.
45. Leith R, O'Connell AC. A clinical study evaluating success of 2 commercially available veneered primary molar stainless steel crowns. *Pediatr Dent.* 2011;33:300–6.
46. Yilmaz Y, Koçoğullari ME. Clinical evaluation of two different methods of stainless steel esthetic crowns. *J Dent Child .* 2004;71:212–4.
47. Beldüz Kara N, Yilmaz Y. Assessment of oral hygiene and periodontal health around posterior primary molars after their restoration with various crown types. *Int J Paediatr Dent.* 2014;24:303–13.
48. Zimmerman JA, Feigal RJ, Till MJ, Hodges JS. Parental attitudes on restorative materials as factors influencing current use in pediatric dentistry. *Pediatr Dent.* 2009;31:63–70.
49. Seale NS, Randall R. The use of stainless steel crowns: a systematic literature review. *Pediatr Dent.* 2015;37:145–60.
50. Bell SJ, Morgan AG, Marshman Z, Rodd HD. Child and parental acceptance of preformed metal crowns. *Eur Arch Paediatr Dent.* 2010;11:218–24.
51. Champagne C, Waggoner W, Ditmyer M, Casamassimo PS, MacLean J. Parental satisfaction with veneered stainless steel crowns for primary anterior teeth. *Pediatr Dent.* 2007;29:465–9.
52. Shah PV, Lee JY, Wright JT. Clinical success and parental

- satisfaction with anterior veneered primary stainless steel crowns. *Pediatr Dent*. 2004;26:391–5.
53. Ashima G, Sarabjot KB, Gauba K, Mittal HC. Zirconia crowns for rehabilitation of decayed primary incisors: An esthetic alternative. *J Clin Pediatr Dent*. 2014;39:18–22.
 54. del Pozo PP, Fuks AB. Zirconia crowns-An esthetic and resistant restorative alternative for ECC affected primary teeth. *J Clin Pediatr Dent*. 2014;38:193–5.
 55. Clark L, Wells MH, Harris EF, Lou J. Comparison of amount of primary tooth reduction required for anterior and posterior zirconia and stainless steel crowns. *Pediatr Dent*. 2016;38:42–6.
 56. Abdulhadi B, Abdullah M, Alaki S, Alamoudi N, Attar M. Clinical evaluation between zirconia crowns and stainless steel crowns in primary molars teeth. *Journal of Pediatric Dentistry*. 2017;38:42–6.
 57. Sharaf AA, Farsi NM. A clinical and radiographic evaluation of stainless steel crowns for primary molars. *Journal of Dentistry*. 2004;32:136–142.
 58. Henderson HZ. Evaluation of the preformed stainless steel crown. *ASDC J Dent Child*. 1973;40:353–8.
 59. Myers DR. A clinical study of the response of the gingival tissue surrounding stainless steel crowns. *ASDC J Dent Child*. 1975;42:281–4.
 60. Checchio LM, Gaskill WF, Carrel R. The relationship between periodontal disease and stainless steel crowns. *ASDC J Dent Child*. 1983;50:205–9.
 61. Myers DR, Schuster GS, Bell RA, Barenie JT, Mitchell R. The effect of polishing technics on surface smoothness and plaque accumulation on stainless steel crowns. *Pediatr Dent*. 1980;2:275–8.
 62. Webber DL. Gingival health following placement of stainless steel crowns. *ASDC J Dent Child*. 1974;41:186–9.
 63. Warren JJ, Yonezu T, Bishara SE. Tooth wear patterns in the deciduous dentition. *Am J Orthod Dentofacial Orthop*. 2002;122:614–8.
 64. Sulong MZ, Aziz RA. Wear of materials used in dentistry: a review of the literature. *J Prosthet Dent*. 1990;63:342–9.
 65. Bimstein E, Zaidenberg R, Soskolne AW. Alveolar bone loss and restorative dentistry in the primary molars. *J Clin Pediatr Dent*. 1996;21:51–4.
 66. Guelmann M, Matsson L, Bimstein E. Periodontal health at first permanent molars adjacent to primary molar stainless steel crowns. *J Clin Periodontol*. 1988;15:531–3.
 67. Vijayashree Priyadharsini J. In silico validation of the non-antibiotic drugs acetaminophen and ibuprofen as antibacterial agents against red complex pathogens. *J Periodontol*. 2019;90:1441–8.
 68. Ezhilarasan D, Apoorva VS, Ashok Vardhan N. Syzygium cumini extract induced reactive oxygen species-mediated apoptosis in human oral squamous carcinoma cells. *J Oral Pathol Med*. 2019;48:115–21.
 69. Ramesh A, Varghese S, Jayakumar ND, Malaiappan S. Comparative estimation of sulfiredoxin levels between chronic periodontitis and healthy patients: A case-control study. *J Periodontol*. 2018;89:1241–8.
 70. Mathew MG, Samuel SR, Soni AJ, Roopa KB. Evaluation of adhesion of *Streptococcus mutans*, plaque accumulation on zirconia and stainless steel crowns, and surrounding gingival inflammation in primary 2020;48:115–21
 71. Sridharan G, Ramani P, Patankar S, Vijayaraghavan R. Evaluation of salivary metabolomics in oral leukoplakia and oral squamous cell carcinoma. *J Oral Pathol Med*. 2019;48:299–306.
 72. Pc J, Marimuthu T, Devadoss P. Prevalence and measurement of anterior loop of the mandibular canal using CBCT: A cross sectional study. *Clin Implant Dent Relat Res*. 2018;89:1241–8.