

# Comparative Evaluation and Correlation of Neurologic Symptoms, Otologic Symptoms and Proximity of TMJ Disk with Condyle in Class II (Vertical) Mild, Moderate and Severe TMD Cases as Compared to Non-TMD Cases

Abhishek Sanchla<sup>1\*</sup>, Sunita Shrivastava<sup>1</sup>, Ranjit Kamble<sup>2</sup>

<sup>1</sup>Department of Orthodontics and Dentofacial Orthopaedics, Datta Meghe Institute of Medical Sciences, Sawangi (Meghe), Wardha (Maharashtra), India; <sup>2</sup>Department of Orthodontics and Dentofacial Orthopaedics, Datta Meghe Institute of Medical Sciences, Sawangi (Meghe), Wardha (Maharashtra), India.

Corresponding author: Abhishek Sanchla, Department of Orthodontics and Dentofacial Orthopaedics, Datta Meghe Institute of Medical Sciences, Sawangi (Meghe), Wardha (Maharashtra), India, Tel: 9930055381; E-mail: abhisheksanchla@gmail.com

## Abstract

**Introduction:** Temporomandibular disorder is a collective term used to describe a number of related disorders affecting the TMJ, muscles of mastication and associated structures all of which have common symptoms like pain and reduced jaw opening. Most of the studies on temporomandibular disorders in literature are related to malocclusion teeth and orthodontic treatment. In literature the role of "Neurologic" and "Otologic" symptoms have been researched in isolations. There is a need to co-relate these findings with the severity of temporomandibular disorders in patients seeking orthodontic treatment. **Objectives:** To evaluate and compare Neurologic symptoms, Otologic symptoms and its correlation with the proximity of TMJ disk and condyle in mild, moderate and severe Class II (vertical) TMD cases. **Methodology:** Grading of TMD using Helkimo and Craniomandibular index and evaluation of Neurologic and Otologic symptoms by separate questionnaires and its correlation with the proximity of TMJ disk and condyle using MRI. **Expected results:** Degree of association of Neurologic and Otologic symptoms with TMD. **Conclusion:** Temporomandibular disorders can be diagnosed by assessing the extra-craniofacial symptoms like Neurologic and Otologic symptoms and may help in early diagnosis and evaluating the potential risk factors for developing temporomandibular disorders. Hence a study is planned in the department.

**Keywords:** Temporomandibular joint disorder; TMD, Helkimo index; craniomandibular index, Neurologic symptoms; Otologic symptoms

## Introduction

Temporomandibular disorder is a broad term used to describe a number of interlinked disorders affecting the TMJ, muscles of mastication and associated structures all of which have common symptoms such as pain and reduced jaw opening. [1,2] According to Rieder et al. the prevalence of patients with at least one symptom of TMD ranged from 33% to 50%.

However the patients that need professional attention range around 10%. [3] The etiology of TMD is complex and multifactorial. There are various factors that can contribute to this disorder such as structural deformities of the joints, pathological alteration of the joints, changes in masticatory musculature, genetic factors, psychological factors micro and macro trauma. [4,5]

According to a study conducted in 2013, Leonor Satnez-prez concluded that there was a correlation between occlusal anomalies, malocclusion and TMD. [6] But there are studies contraindicating this and stating that there is no direct correlation between malocclusion and TMD (Turp J Schindler, 2012). [7] Various other studies have been documented in literature relating TMD and orthodontic treatment, most of them concluding a negative co-relation (Mc. Namra 1995). [8]

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 3.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as the author is credited and the new creations are licensed under the identical terms.

**How to cite this article:** Sanchla A, Shrivastava S, Kamble R. Comparative Evaluation and Correlation of Neurologic Symptoms, Otologic Symptoms and Proximity of TMJ Disk with Condyle in Class II (Vertical) Mild, Moderate and Severe TMD Cases as Compared to Non-TMD Cases. AMHSR. 2021;11:102-105

Most of the studies in literature are related to malocclusion teeth and orthodontic treatment and these extensive studies have led to the development of diagnostic criteria for TMD. [9]

As Orthodontists we have studied the above association but there are structure other than the craniofacial structures, that may also help in easy detection of temporomandibular disorders in patients seeking orthodontic treatment that needs to be researched and co-related as they may be beneficial in taking measures and stop the future progress of the condition. [10] In literature the role of “Neurologic” and “Otologic” symptoms have been researched in isolations. Also the proximity of the mandibular nerve and ear to the condyle and articular disc of TMJ may have a role in the symptomatology of TMD. [11] There is a need to co-relate these findings with the severity of temporomandibular disorders in patients seeking orthodontic treatment. Hence a study is planned in the department.

## Aim and Objectives

### Aim

To evaluate ,compare and co-relate Neurologic symptoms, Otologic symptoms and proximity of TMJ disk and condyle in class II (vertical) mild, moderate and severe temporomandibular cases as compared to non-TMD cases

### Objectives

- 1.Grading of TMD using Helkimo index.
- 2.To evaluate Neurologic and Otologic symptoms in Class II(vertical) mild, moderate and severe temporomandibular disorder cases.
- 3.To evaluate the proximity of TMJ disk and condyle in Class II(vertical) mild, moderate and severe temporomandibular disorder cases.
- 4.To compare Neurologic and Otologic symptoms in Class II(vertical) mild, moderate and severe TMD cases with non-TMD cases.
- 5.To compare the proximity of TMJ disk and condyle in Class II (vertical) mild, moderate and severe TMD cases with non-TMD cases.
- 6.To correlate the Neurologic and Otologic symptoms with the proximity of TMJ disk and condyle in Class II (vertical) mild ,moderate and severe TMD cases.

### Study design

#### Inclusion criteria

- 1.Patients with Class I malocclusion (diagnosed using cephalometric analysis)
- 2.Patients with Class II (vertical) malocclusion and temporomandibular disorder

3.All permanent teeth present

4.Adult age group [Table 1]


#### Exclusion criteria

- 1.Non –TMD class II(vertical ) cases.
- 2.All Class II (horizontal) cases.
- 3.Class III malocclusion cases.
- 4.Patients with myofascial pain dysfunction and myalgia.
- 5.Previous intervention with TMJ surgery or trauma to TMJ
- 6.Patients with bony disorders.
- 7.Previous with history of Neurological disorders.
- 8.Patients with gross pathology of ears

### Study methodology

1. The observational study will be conducted in the Department Of Orthodontics And Dentofacial Orthopaedics ,Sharad Pawar Dental College, Sawangi(M),Wardha. Approval from the Ethical Committee has been obtained.

2. Total of 60 adult patients (class I, class II vertical) will be selected from the Outpatient Department(OPD) of Orthodontics and Dentofacial Orthopaedics of Sharad Pawar Dental College, Sawangi, Wardha [Figure 1].



**Figure 1:** Flow Chart of adult patient study.

For observing all universal precautions and infection control procedures will be taken. The patients will be divided into 4 groups:

Group A(control group)-15 Class I non-TMD cases.

Group B-15 Class II (vertical) mild TMD cases.

Group C-15 Class II (vertical) moderate TMD cases.

Group D-15 Class II (vertical) severe TMD cases.

1.Informed and written consent will be obtained from all the patients. For all the patients.

2. Helkimo anamnesic and clinical dysfunction indices will be obtained.
3. Craniomandibular index will be obtained for all patients
4. Based on this temporomandibular disorders will be graded into mild moderate and severe cases.
5. The Neurologic and Otologic symptoms will be evaluated by separate questionnaires.
6. The proximity of the TMJ disk and condyle will be assessed using MRI and will be co-related with the outcomes of the questionnaire evaluation of Neurologic and Otologic symptoms.

### Sample Size

The calculation of sample size was calculated based on the values from the article Comparison of the efficacy of a questionnaire, oral history, and clinical examination in detecting signs and symptoms of occlusal and temporomandibular joint dysfunction

$$n = N * X / (X + N - 1)$$

Considering the prevalence of severe cases as 0.95%, 15 patients would be taken in each group.

15 patients in 4 groups Hence , total sample size=60

### Expected Results

The Neurologic and Otologic symptoms may be associated with temporomandibular disorders depending on the severity of the disorder and there may be a correlation between these symptoms and the proximity of TMJ disk and condyle.

### Discussion

The incidence of temporomandibular disorders is constantly increasing. Also the causative factors of TMD is said to be of multiple origin which has led to the formulation of diagnostic criteria for TMD. But all the symptoms assessed for its diagnosis are related to symptoms of craniofacial structures like malocclusion, habits or bone deformities. Assessing the Neurologic and Otologic symptoms in patients with mild, moderate and severe cases of TMD will lead to a broad spectrum approach for diagnosis and prevention of risk factors that lead to development of temporomandibular disorders. Pamela Maria Kusda conducted an analytical study relating Otological symptoms with temporomandibular disorders in male and female patients, TMD evaluated by RDC/TMD. Records of 485 patients were analysed for prevalence of ear complaints. The study concluded that Otological symptoms like tinnitus, deafness, fullness in ears ,dizziness were associated with TMD regardless of age and sex. [12] E. Pedulla evaluating the cause of neuropathic pain in patients with temporomandibular disorders using Magnetic Resonance Imaging (MRI). A total of 48 patients (16 with TMD and neuropathic pain, 16 with TMD and no neuropathic pain and 16 healthy patients) were evaluated. The distance between mandibular nerve and TMJ disk was

evaluated using MRI. The study concluded that neuropathic pain in patients with TMD may be due to the close proximity between TMJ disk and mandibular nerve. [13] Other related studies by Helkimo, [14] Friction, [15] Awasthi. [16] Sundrani [17], Muraraka [18] were reviewed. Gupta reported about stress distribution in the temporomandibular joint after mandibular protraction [19]. Similar study was reported by Shrivastava [20].

### Conclusion

No such studies has been carried out to evaluate the extra-craniofacial symptoms in temporomandibular disorders correlating it to TMJ disk proximity. This may help in early diagnosis of TMD and evaluating the potential risk factors for developing temporomandibular disorders and hence avoiding or delaying its development.

### Scope

This study will help us to diagnose TMD's by assessing the extra-craniofacial symptoms (Neurologic and Otologic symptoms). An early diagnosis will be possible and will thus help in decreasing the severity of TMD or avoiding its development.

### Reference

1. Dimitroulis G. Temporomandibular disorders: a clinical update. *Bmj*. 1998;317:190-4.
2. John ZA, Shrivastav SS, Kamble R, Jaiswal E, Dhande R. Three-dimensional comparative evaluation of articular disc position and other temporomandibular joint morphology in Class II horizontal and vertical cases with Class I malocclusion: A magnetic resonance imaging study. *Angle Orthod*. 2020;90:707-14.
3. Rieder CE. Comparison of the efficacy of a questionnaire, oral history, and clinical examination in detecting signs and symptoms of occlusal and temporomandibular joint dysfunction. *J Prosthet Dent*. 1977;38:433-40.
4. Gupta R, Luthra RP, Kaur D, Aggarwal B. Temporomandibular disorders: A review. *Int. J. Adv. Sci. Res*. 209;4:22-26.
5. GOSAVI DS, SHRIVASTAV DS. Comparative Analysis For The Presence And Intensity Of TMD Symptoms In Skeletal Class I Malocclusion, Skeletal Class II Horizontal Malocclusion & Skeletal Class II Vertical Malocclusion Using Helkimo And Craniomandibular Index. A Study Protocol. *EJMCM*. 2020;7:2113-8.
6. Sánchez-Pérez L, Irigoyen-Camacho ME, Molina-Frechero N, Mendoza-Roaf P, Medina-Solís C, Acosta-Gío E, Maupomé G. Malocclusion and TMJ disorders in teenagers from private and public schools in Mexico City. *Med Oral Patol Oral Cir Bucal*. 2013;18:312-318.
7. Türp JC, Schindler H. The dental occlusion as a suspected cause for TMDs: epidemiological and etiological considerations. *J. Oral Rehabil*. 2012;39:502-12.

8. McNamara Jr JA. Orthodontic treatment and temporomandibular disorders. *Oral Surgery. Oral Surg Oral Med Oral Pathol Oral Radiol Endod.* 1997;83:107-17.
9. Dhannawat D, Purva V, Shrivastav D, Kamble D, Banerjee D. Different Types of Occlusal Splint Used In Management of Temporomandibular Joint Disorders-A Review. *EJMCM.* 2020;7:1787-94.
10. John ZA, Shrivastav SS, Kamble R, Jaiswal E, Dhande R. Three-dimensional comparative evaluation of articular disc position and other temporomandibular joint morphology in Class II horizontal and vertical cases with Class I malocclusion: A magnetic resonance imaging study. *Angle Orthod.* 2020;90:707-14.
11. Bouhassira D, Attal N, Alchaar H, Boureau F, Brochet B, Bruxelle J, Cunin G, Fermanian J, Ginies P, Grun-Overdyking A, Jafari-Schluep H. Comparison of pain syndromes associated with nervous or somatic lesions and development of a new neuropathic pain diagnostic questionnaire (DN4). *Pain.* 2005;114:29-36.
12. Kusdra PM, Stechman-Neto J, de Leão BL, Martins PF, de Lacerda AB, Zeigelboim BS. Relationship between Otological symptoms and TMD. *Int. Tinnitus J.* 2018;22:30-4.
13. Cascone P, Di Paolo C, Leonardi R, Pedullà E. Temporomandibular disorders and orthognathic surgery. *J CRANIOFAC SURG.* 2008;19:687-92.
14. HELKIMO M. Studies on function and dysfunction of the masticatory system. III. Analysis of anamnestic and clinical recording of dysfunction with the aid of indices. *Swed Dent J.* 1974;67:165-73.
15. Friction JR, Schiffman EL. The craniomandibular index: validity. *J Prosthet Dent.* 1987;58:222-8.
16. Awasthi E, Bhola N, Kamble R, Shrivastav S and Goyal A. Treatment of an Adult with Skeletal Class III and A Hemimandibular Elongation A Multidisciplinary Approach. *J. Clin. Diagnostic Res.* 2016;10:5-6.
17. Sundrani A, Kamble RH, Ahuja MM, Bhola N and Shrivastav S. Surgical Orthodontic Correction of Class II Malocclusion with Vertical Maxillary Excess and Gummy Smile- A Case Report. *J. Evol. Med. Dent. Sci.* 2020;9:67-70.
18. Murarka SP, Shrivastav S, Kamble R, Dargahwala H, Khakhar P, John Z, et al. Comparative Evaluation of Discomfort, Expectations and Functional Experiences during Treatment of Class II Malocclusion with Forsus Fixed Functional Appliance and Sharma's Class II Corrector - A Questionnaire Based Survey. *J. Evol. Med. Dent. Sci.* 2021;10:474-78.
19. Gupta A, Hazarey PV, Kharbanda OmP, Kohli VS and Gunjal A. Stress Distribution in the Temporomandibular Joint after Mandibular Protraction: A 3-Dimensional Finite Element Study. Part 2. *Am. J. Orthod. Dentofac. Orthop.* 2009;135:749-56.
20. Shrivastava, Abhinav, Hazarey PV, Kharbanda OmP and Gupta A. Stress Distribution in the Temporomandibular Joint after Mandibular Protraction: A Three-Dimensional Finite Element Study. *Angle Orthod.* 2015; 85:196-205.