

Nigeria and the Selenium Micronutrient: A Review

Adeniyi MJ* and Agoreyo FO

Department of Physiology, University of Benin, Benin-city, Nigeria

Corresponding author:

Adeniyi MJ,
Department of Physiology, University
of Benin, Benin-city, Nigeria,
Tel: +23408066796517;
E-mail: 7jimade@gmail.com

Abstract

Local studies have documented the contributions of minerals to the health status of mammals. Selenium (Se) is an example of such minerals. It is essential for the activation and functioning of enzymes. In Nigeria, the mineral has been reported to be present in water, foods and soils. While the soil selenium concentrations determine selenium levels in foods, difference in food concentrations of the mineral atones for the variations in the plasma and tissue selenium level across the geopolitical zones of Nigeria. Areas in Nigeria prone to erosion could be associated with inadequate soil selenium levels and this might lead to low availability of the mineral in the foods consumed by animals and humans. The plasma selenium levels in Nigerians may vary from as low as 0.006 mg/L to as high as 0.3279+/-0.0546 mg/L depending on the geographical factors and analytical methods employed. Studies in which the physiological effects of selenium supplementations were investigated revealed an improvement in CD4 counts in HIV positive human subjects, reduction in rats' susceptibility to trypanosomiasis, improvement in reproductive function, acceleration of ulcer healing in animal model of experimental gastric ulcer and mitigation of the adverse effects in animal model of gastric ulcer. Low plasma selenium levels are associated with HIV infection, fertility problems, postpartum cardiomyopathy and diabetes mellitus. Therefore, there is growing need for dietary selenium supplementations in Nigeria.

Keywords: Selenium; Supplementations; Nigeria; Geopolitical; Nigerian; Mineral

Introduction

Nutrition deals with food substances and their effects on health. [1] Food substances including non-energy yielding substances like selenium are required for metabolisms. [2] Selenium is an element of the periodic table and it is present in soil, water, rock and food. As far as Nigeria is concerned, studies revealed that the soil concentrations of the mineral vary across the six geopolitical zones of the country. This results in variation in concentration of the mineral in plant and animal foods consumed by Nigerians [3,4] and difference in selenium level in body fluid and tissues. [5,6]

As a vital component of antioxidant enzymes [7] and iodothyronine deiodinase, [8] the mineral mediates many functions in humans and animals. For instance, selenium improved glutathione peroxidase and CD4 count in HIV positive Nigerian patients. [9-11] It also exerted a positive influence on male reproductive function [12] and accelerated gastric ulcer healing in an animal study. The mineral is essential for cardiovascular health [13] and endocrine functions. [14] Selenium has also been shown to possess insulinotropic, [15] antioxidant [7] and anticancer effects. [16] Selenium, among other functions, reduced susceptibility to infection in rats. [17]

The purpose of this work is to highlight trends in selenium research and the physiological significance of its supplementations in Nigeria [Figure 1].

Literature Review - Selenium

Chemistry of selenium

Selenium is a metalloid located in period IV and group VI of the periodic table. It has atomic number 34 and mass number

of 79 with oxidation states of +6, +5, +4, +3, +2, +1 and -1. [18] It rarely occurs in its elemental state in nature, or as pure ore compounds. [18] Selenium forms two oxides: selenium dioxide and selenium trioxide. Selenium dioxide dissolves in water to form selenous acid. Salts of selenous acid are called selenites and these include silver selenite and sodium selenite, the inorganic forms of selenium. Selenium forms hydrogen selenide (H₂Se) which is toxic and colourless. Selenium forms stable bonds to carbon giving rise into selenols. Examples of selenols are diphenyldiselenide, benzeneselenol and selenomethionine, the organic forms of selenium. [18]




Figure 1: Map of Nigeria (Perry-Castaneda Library, 101 East 21st St.)

The mineral was first discovered in 1817 by Jons Jacob Berzelius when investigating the chemicals responsible for outbreaks of ill health among workers in a Swedish sulphuric

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 3.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as the author is credited and the new creations are licensed under the identical terms.

How to Cite this Article: Adeniyi MJ, et al. Nigeria and the Selenium Micronutrient: A Review. *Ann Med Health Sci Res.* 2018;8:5-11

states, South-South geopolitical zone of Nigeria was 2.86 ± 3.97 mg/dl.^[40] Factors which may account for variation of selenium in Nigeria are:

Table 1: Mammalian selenoproteins.^[21]

S/n	Selenoprotein	Proposed function
	GPx	
1	GPx1	Antioxidant In cell cytosol; Se store
2	GPx2	Antioxidant In gastrointestinal tract
3	GPx3	Antioxidant In extracellular space and plasma
4	GPx4	Antioxidant In membranes, structural protein in sperm: apoptosis
5	GPx5	Unknown
6	GPx6	GPx1 homolog?
		Multiple roles including dithiol-disulphide oxidoreductase
7	Thioredoxin reductase (TR)	Detoxifies peroxides, reduces thioredoxin (control of cell growth): maintains redox state of transcription factors
8	TR1	Mainly cytosolic, ubiquitous
9	TR2	Expressed by testes
10	TR3	Mitochondrial, ubiquitous
11	Iodothyronine deiodinases	
12	Types D1 AND D2	Converts thyroxine (T4) to bioactive 3,6,3'-tri-iodothyronine (T3)
13	Types D1 and D3	Converts T4 to to inactive 3',3',5' reverse T3
14	Salenoprotein P	Selenium transport protein, antioxidant in endothelium
15	Salenoprotein W	Antioxidants in cardiac and skeletal muscle
16	Salenophosphate synthetase (SPS2) 15 kDa	Synthesis of selenophosphate for selenoprotein synthesis
17	selenoprotein (sep 15)	Protects against cancer
18	Selenoproteins H, I, K, M, N, O, P, R, S, T, V	Role largely unknown

- **Rainfall:** In Nigeria, the coastal region around Niger Delta region in South-Southern region of Nigeria receives annual around 4,000 mm (157.5 in) of rainfall. In North Eastern state, Borno, the annual rainfall is below 1500mm. In most Southern States, the annual rainfall is above 2,000 mm. Selenium tends to be concentrated in the soil of the drier regions of the world.^[41]

- **Soil pH:** In acidic poorly aerated soils, selenium is relatively unavailable to plants because it is present mainly as insoluble selenite complexes.^[41]

- **Geographical factors:** The selenium content of food varies widely depending on geographic locations.^[42] For example, Nanka community of Anambra state in South-Eastern Nigeria, is an erosion ridden environment. Soil selenium in this area was reported to be low.^[6] Low soil selenium correlates positively with secondary diseases. An example of secondary diseases is Keshan disease, a disease that affects some people that live in South-West China, where the soil is selenium deficient.

- **Soil composition:** Phosphates present in rock, shales, limestones and sand stone contain selenium. Sulphide mineral deposit reduces selenium level.

- **Analytic methods:** Methods devised in estimating soil selenium level give results which may not be the same.^[5,43]

Selenium concentration in Nigerian foods

Since plants depend on raw materials of soil origin, soil selenium content determines selenium availability in the body. Exposure to selenium is majorly through food. Brazilian nut has the highest selenium concentration 0.54 mg/kg.

In Anambra state, South-Eastern region, analysis of cassava tuber in Nanka and Oba communities showed that selenium contents were low: 0.16+/-0.03 ppm, 0.15+/-0.02 ppm respectively.^[6] Tubers are a major staple food in Nigeria, especially in rural communities. Decreased selenium in tubers may portend risk to the health of people [Table 2].

Table 2: Selenium concentrations in Nigerian foods.^[5]

Sample	Mean \pm SD
Cereals	
Millet (<i>Sorghum vulgare</i>)	5 \pm 3.4
Rice (white) (<i>Zea mays</i>)	5 \pm 2.5
Meat and fish	
Fish (dry) (<i>Tilapia nilotica</i>)	15 \pm 4.5
Cray fish (<i>Procambaris clarkia</i>)	20 \pm 5.2
Snail (<i>Achatina fulica</i>)	<2 \pm 3.2
Wistar strain (albino rat)	<1 \pm 8.4
Milk and dairy product	
Milk (powder)	5 \pm 3.5
Egg (yolk)	<1 \pm 8.4
Egg (white)	<2 \pm 3.2
Vegetables	
Water leaf (<i>Talinum triangulare</i>)	5 \pm 3.5
Spinach (<i>Amaranthus spp</i>)	<1 \pm 8.4
Bitter leaf (<i>Vernonia amygdalina</i>)	<1 \pm 8.4
Onion (<i>Allium cepa</i>)	<2 \pm 3.2
Pepper (<i>Capsicum annum</i>)	<2 \pm 3.2

Study by Abulude et al.^[5] examined both local and imported products. It is interesting to observe that cassava flour, Cray fish and tilapia had the highest selenium concentrations. Wistar rats, snails, egg yolk, egg white, spinach, pepper and bitter leaf contained low selenium levels. Conversely, in Northern Europe, foods which contained high selenium level include meat, poultry, and fish (49). Difference in soil selenium levels, presence of high level of phytate, sulphur and mercury and difference in analytical methods employed might account for the discrepancy between the low selenium level reported by Olife and Anajekwu^[6] and the findings of Abulude et al.^[5]

Analysis of average selenium content in root and tuber plants in North-Central by Zarmai et al.^[43] showed that sweet potato contains 19.2+/- 5.20 μ g/kg, yellow yam has 18.3+/-6.97 μ g/kg, yam has 13.6+/-7.12 μ g/kg and cassava has 13.0+/-5.84 μ g/kg. Yellow yam from Lafia had the highest level 76.3 μ g/kg while Minna, Bida, Keffi, Makurdi, Otukpo, Lokoja and Ilorin had a mean of 18.3+/-6.97 μ g/kg. Sweet potato from Bida showed the highest selenium level while Makurdi, Lafiagi and Jos had a mean value of 19.2+/-5.20 μ g/kg. Yam from Makurdi contained the highest selenium concentration 81.8 μ g/kg while that from Bida, Abuja, Lafia, Gboko, Lokoja, Idah, Ilorin, Lafiagi and Jos showed a mean of 13.6+7.12 μ g/kg.

From the work of Zarmai et al.^[43] sweet potato in North-Central region had the highest selenium content followed by yellow yam, yam and cassava.

Study by Orisakwe et al.^[44] estimated the levels of selenium in fifty (50) beverages purchased in Nigeria using the Atomic Absorption Spectrophotometer (AAS). The result indicated that selenium levels ranged from 0.24 - 1.67 mg/L for canned beverages and 0.07 - 1.23 mg/L for non-canned beverages. 95% of canned beverages had selenium levels that exceeded the maximum Contaminant Level (MCL) set by United States Environmental Protection Agency (USEPA) (MCL) whereas 90% of the non-canned products had selenium levels above the MCL [Table 3].

Table 3: Selenium content in North-Central region of Nigeria.^[43]

Tuber	City with highest selenium level	City with lowest selenium level
Yam	Makurdi	Bida, Abuja, Lafiagi and Jos
Sweet potato	Bida	Makurdi, Lafiagi and Jos
Yellow yam	Lafiagi	Minna, Bida, Keffi, Makurdi, Otukpo, Lokoja and Ilorin

Different brands of bottled beer marketed in Makurdi town exhibited selenium content that exceeded maximum contaminant level. On the contrary, Obahiagbon et al.^[45] reported that selenium concentration in vegetable, waterleaf, ewedu leaf, bitter leaf, cocoyam leaf, green pepper leaf and okra leaf were within permissive level.

The above findings showed the variations in food selenium levels. Local foods grown in erosion prone regions may have lower selenium concentrations. This may lead to a decrease in selenium level in animals and humans that consume the foods.

Human selenium requirement

The recommended daily allowances (RDA) of selenium at 55 µg/day (WHO/FAO/IAEA, 1996) for both adult males and females by Food and Nutrition Board at the Institute of Medicine. This amount is based on the amount of dietary selenium needed for activity of selenium-containing enzyme glutathione peroxidase in the plasma. The tolerable upper limit of selenium is 0.4mg/day (WHO/FAO/IAEA, 1996). Excess consumption of selenium may result in brittle nail, garlic breath, nausea and vomiting. 10 gram oral dose of selenium dioxide can cause death. The lethal doses (LD50) for orally administered selenate in rats is 7 to 22 mg/kg body weight^[46] and between 3 and 12 mg/kg body weight for orally administered sodium selenite in rats.^[47,48]

Selenium levels in Nigerians

Kolawole and Obueh^[40] reported that the mean serum selenium level of people in selected areas of Akwa Ibom and Cross River state, South-South region of Nigeria, was 0.006 mg/L. Okwara et al.^[49] got 0.29±/− 0.09 mmol/L (57.6±/− 14.4mg/L) for healthy male adults and 0.28±/−0.08 mmol/L (50.4±/−16.2 mg/L) for healthy adult female in Orlu, South-East Nigeria. Nwagha et al.^[50] also found a plasma selenium concentration of 1.38±/−0.18 µmol/l (0.245±/−0.0324 mg/L) in healthy female subjects in the South-East region.

The mean plasma selenium of 88 healthy adults in Abeokuta, South-Western region of Nigeria, was 0.188±/−0.026 mg/L.^[51] Arinola et al.^[52] obtained a level of 57.05±/−2.50 µg/L (0.057

±/− 0.0025 mg/L) in healthy female Nigerians in South-Western region. The plasma selenium level in a study by Anyabolu et al.^[10] was 9.81±/−1.6 µg/dL (0.0981±/−0.016 mg/L) in children in South-Western region. Sixty healthy subjects of over 18 years of age in an area in Osun state, South-Western region, had a mean serum selenium level of 32.79±/−5.46 µg/dl (0.3279±/−0.0546 mg/L).^[53] Oyedeji et al.^[54] also obtained a plasma level of 1.02±/−0.01 µmol/L (0.1836±/−0.0018 mg/L) in healthy female subjects in South-Western region. Karaye et al.^[13] obtained a plasma level of 118.4 ± 45.6 µg/L (0.118±/−0.0456 mg/L) in a selected area of North-West region of Nigeria.

In addition to plasma selenium level, physiological balance in selenium distribution between body fluid compartments may be important as far as state of health is concerned. For instance, the ratio of serum selenium to seminal plasma selenium was found to be 1:1 in healthy adult male in South-West of Nigeria.^[12]

In developing country like Nigeria where a sizeable population of people lives on local foods which may not contain high selenium contents due to soil factors, the selenium concentrations in body fluids would be expected to vary across the geopolitical zones. Despite this, the selenium content in Nigeria putting all these studies together may vary from 0.006 mg/L to 0.3279±/−0.0546 mg/L.

Factors which may affect the plasma and tissue concentrations of the mineral may interfere with the physiological functions of the mineral. Such factors may include:

- Age: Plasma selenium is lower in children than adult.^[55]
- Genetic variation affects the plasma and tissue levels of selenium.^[56]
- Gender: Males have slightly higher plasma selenium than female.^[55]
- Pathological conditions including Acquired Immune Deficiency Syndrome,^[43,57,58] infertility,^[12] diabetes mellitus^[14] and digestive problems^[12] decrease plasma and tissue selenium.
- Pregnancy and lactation^[50] decrease plasma selenium level.
- State of activity: Exercise increases the need for selenium^[32] and sedation may reduce plasma selenium level.^[12]
- Kidney dialysis reduces plasma selenium.^[12]
- Lifestyle such as smoking, alcoholism and vegetarianism may reduce plasma selenium level.^[55]
- Oral contraceptive use^[51,59] reduces selenium level.

Physiological Functions of Selenium

Immunity

Many local studies have documented that HIV-positive subjects exhibited low plasma selenium with male subjects having lower serum selenium levels than female subjects.^[9-11,60] Report also indicates that HIV positive subjects with and without skin diseases exhibited low serum selenium.^[61] The virus is known to incorporate selenium into its Human-Like glutathione peroxidase leading to a decline in serum selenium and glutathione in the host.^[62]

The prevalence of HIV infected children is high.^[10] Therefore, there is incontestable need for selenium supplementation. Selenium is an active component of selenoproteins which include thioredoxin reductase and glutathione peroxidase. In Nigeria, the plasma glutathione was estimated in a study as 0.127±0.022 U/mL. Therefore, selenium supplementation may improve glutathione defense system. Positive immune outcome achieved with respect to selenium supplementation includes improvement in CD4 counts in HIV-positive Nigerian subjects.^[9] In West Africa, Senegal has the highest soil selenium contents and this has been linked with lower prevalence of HIV in the country.^[35]

In animal studies, selenium supplementation together with vitamin E, increased packed cell volume in trypanosome *Brucei* infected rats.^[17] It also reduced susceptibility of black marshal cockerel to infectious bursal disease.^[63]

Reproduction

Selenium supplementation may improve testosterone level in male. Testes and ovaries are among organs that contain greatest concentrations of free selenium.^[25] Upset in physiological balance between serum and seminal plasma selenium was associated with oligospermia and azoospermia in a study by Akinloye et al.^[12] Selenium supplementation improved reproductive of rabbits does by enhancing follicle stimulating hormone and oxidative enzyme. Low selenium level was found in subjects with prostate cancer.^[58]

Endocrinology and metabolism

Female diabetics drawn from Calabar showed higher serum selenium level than diabetic male.^[14] Together with zinc, selenium reduced blood glucose concentration, restored hepatic functions and reduced hepatic and renal lipid peroxidation of diabetic rats.^[64] Ahmed et al.^[15] reported that selenium possessed a hypoglycemic property comparable to glibenclamide in rats.

Gastrointestinal health

Selenium supplementation may also exert positive influence on gastrointestinal function and health. In an animal experiment conducted by Adeniyi et al. supplementation with selenium accelerated gastric ulcer healing by facilitating mucosal regeneration and antioxidant activity and by reducing lipid peroxidation. Together with vitamin E, selenium demonstrated tendency to assuage adverse effects in an experimental model of acute gastric ulceration in rats.^[65]

Cardiovascular health

In South western China, selenium deficiency was associated with cardiomyopathy called Keshan disease. Also in Nigeria, a study by Karaye et al. in an area in North-West region of Nigeria indicated that female subjects with Postpartum Cardiomyopathy had low serum selenium level that was due to insufficient dietary intake of the mineral.

Conclusion

There are variations in selenium levels across the geopolitical zones in Nigeria with a nadir of 0.006mg/L and a zenith of 0.3279±0.0546mg/L. Hence, there is need for selenium

supplementation. This review also provides evidence that selenium supplementations can improve health status of Nigerians.

Conflict of Interest

All authors disclose that there was no conflict of interest.

References

1. Ellie W, Rady RS. Understanding Nutrition (13th ed.), Wadsworth. 2013;667-670.
2. Opstad K. Circadian rhythm of hormones is extinguished during prolonged physical stress, sleep and energy deficiency in young men. *Eur J Endocrinol.* 1994;131:56-66.
3. Rayman MP. The importance of selenium to human health. *Lancet.* 2000;356:233-241.
4. Combs GF Jr. Selenium in global food systems. *Br J Nutr.* 2001;85:517-547.
5. Abulude F, Ogunkoya M, Orojo T. Selenium In Nigerian Foods. *Ejipau Anim. Reprod.* 2006;10:677-683.
6. Ifeyinwa OC, Azuka AB. Iodine and selenium Distribution in The Local Environment of Selected Villages in Anambra State, Nigeria. *Journal of Natural Sciences Research.* 2013;3:116.
7. Rayman MP. Selenium and human health. *Lancet.* 2012;379:1256-1268.
8. Mistry HD, Pipkin FB, Redman CW, Poston L. Selenium in reproductive health. *American Journal of Obstetrics & Gynecology.* 2012;206:21-30.
9. Odunukwe NN, Onwujekwe D, Ezechi OO, Ezobi P, Gbajabiamila T, Anyanwu R, et al. The role of selenium as adjunct to HAART among HIV-infected individuals who are advanced in their disease. In Sixteenth International AIDS Conference, Toronto, Canada 2006.
10. Anyabolu HC, Adejuyigbe EA, Adeodu OO. Serum micronutrient status of Haart-naive, HIV infected children in South Western Nigeria: A case controlled study. *AIDS research and treatment.* 2014;2014.
11. Kalio IS. Comparative study of selenium and zinc antioxidant in HIV sero positive individuals.
12. Akinloye O, Arowojolu AO, Shittu OB, Adejuwon CA, Osotimehin B. Selenium status of idiopathic infertile Nigerian males. *Biological trace element research.* 2005;104:9-18.
13. Karaye KM, Yahaya IA, Lindmark K, Henein MY. Serum selenium and ceruloplasmin in Nigerians with peripartum cardiomyopathy. *International Journal of Molecular Sciences.* 2015;16:7644-7654.
14. Nsonwu AC, Usoro CO, Etukudo MH, Usoro IN. Influence of age, gender and duration of diabetes on serum and urine levels of zinc, magnesium, selenium and chromium in type 2 diabetics in Calabar, Nigeria. *Turk J Biochem.* 2006;31:107-114.
15. Ahmed MK, Aliyu M, Yusuf T, Yusuf MK. Ameliorative effect of selenium yeast on blood glucose level in streptozotocin induced diabetes in wistar rats. *Drugs.* 1996;80:23.
16. Zhang X, Zarbl H. Chemopreventive doses of methylselenocysteine alter circadian rhythm in rat mammary tissue. *Cancer Prevention Research.* 2008;1:119-27.
17. Yakubu DP, Dawet A, Olaleye NA. Effects of vitamin E and selenium on some blood parameter of *Trypanosoma brucei brucei* infected rats. *British Journal of Applied Science & Technology.* 2014;4:1100-1108.
18. Greenwood NN, Earnshaw A. *Chemistry of the Elements* (2nd ed). Butterworth-Jheinemann. 1997;751-752.
19. Oldfield JE. The two faces of selenium. *The Journal of Nutrition.* 1987;117:2002-2008.
20. Schwarz K, Foltz CM. Selenium as an integral part of factor 3 against dietary necrotic liver degeneration. *Journal of the American Chemical Society.* 1957;79:3292-3293.
21. Reilly C. *Selenium in food and health.* Springer Science & Business Media; 1996.

22. Robinson MF, Godfrey PJ, Thomson CD, Rea HM, Van Rij AM. Blood selenium and glutathione peroxidase activity in normal subjects and in surgical patients with and without cancer in New Zealand. *The American Journal of Clinical Nutrition*. 1979;32:1477-1485.
23. Kryukov GV, Castellano S, Novoselov SV, Lobanov AV, Zehntab O, Guigó R, et al. Characterization of mammalian selenoproteomes. *Science*. 2003;300:1439-1443.
24. Thomson CD, Stewart RD. Metabolic studies of [75 Se] selenomethionine and [75 Se] selenite in the rat. *British Journal of Nutrition*. 1973;30:139-147.
25. Drasch G, Mailänder S, Schlosser C, Roeder G. Content of non-mercury-associated selenium in human tissues. *Biological trace element research*. 2000;77:219-230.
26. Kohrle J, Jakob F, Contempre B, Dumont JE. Selenium, the thyroid, and the endocrine system. *Endocrine reviews*. 2005;26:944-984.
27. Sunde RA. Molecular biology of selenoproteins. *Annual Review of Nutrition*. 1990;10:451-474.
28. Ip C. Lessons from basic research in selenium and cancer prevention. *The Journal of nutrition*. 1998;128:1845-1854.
29. Robinson MF, McKenzie JM, Thompson CD, Van Rij AL. Metabolic balance of zinc, copper, cadmium, iron, molybdenum and selenium in young New Zealand women. *British journal of nutrition*. 1973;30:195-205.
30. Oster O, Prellwitz W. The renal excretion of selenium. *Biol Trace Elem Res*. 1990;24:119-146.
31. Krupinski T, Beitel GJ. Unexpected roles of the Na-K-ATPase and other ion transporters in cell junctions and tubulogenesis. *Physiology*. 2009;24:192-201.
32. Hurst R, Armah CN, Dainty JR, Hart DJ, Teucher B, Goldson AJ, et al. Establishing optimal selenium status: results of a randomized, double-blind, placebo-controlled trial-. *The American Journal of Clinical Nutrition*. 2010;91:923-931.
33. Okpo U, Christian EC, Nnaemeka E. *Global Journal of Science*. Frontier Research. 2013.
34. National Toxicology Program Report. NTP Technical report on toxicity studies of sodium selenate and sodium selenite administered in drinking water to F344/N rats and B6C3F1 Mice. NC 27709 NIH. 1994.
35. Lawrence W. Selenium and its mighty benefits. *Journal Alternatives*. 2004;12:105-111.
36. Melse-Boonstra A, Hogenkamp P, Lungu OI. Mitigating HIV/AIDS in Sub-Saharan Africa through selenium in food. Farmer Publication, Lusaka. Golden Valley Agricultural Research Trust (GART). 2007;11.
37. Aremu MO, Atolaiye BO, Labaran L. Environmental implication of metal concentrations in soil, plant foods and pond in area around the Derelict Udege mines of Nasarawa State, Nigeria. *Bulletin of the Chemical Society of Ethiopia*. 2010;24.
38. Raji MI, Ibrahim YK, Ehinmidu JO. Physico-chemical characteristics and Heavy metal levels in Drinking Water sources in Sokoto metropolis in North-western Nigeria. *Journal of Applied Sciences and Environmental Management*. 2010;14.
39. Oklo AD, Sha'Ato R, Asemave K, Mkurzurum C. Selenium Content in some Brands of Bottled Beer Marketed in Makurdi Town Benue State, Nigeria. *Int. J. Sci. Res*. 2013;2:457-459.
40. Kolawole SE, Obueh HO. Relationship between soil contents and plasma levels of selenium, chromium and manganese in healthy adult Nigerians. *African Journal of Biotechnology*. 2013;12.
41. Stangoulis J, Graham R, Lyons G. Nutri-prevention of disease with high selenium wheat. *Journal of the Australasian College of Nutritional and Environmental Medicine*. 2003;22:3.
42. Rayman MP. Food-chain selenium and human health: emphasis on intake. *British journal of nutrition*. 2008;100:254-268.
43. Zarmai S, Eneji IS, Sha'Ato R. Analysis of selenium content in root and tuber plants in central Nigeria. *American Journal of Analytical Chemistry*. 2013;4:739.
44. Orisakwe OE, Oragwu CI, Maduabuchi JM, Nzegwu CN, Nduka JK. Copper, selenium and zinc content of canned and noncanned beverages in Nigeria. *African Journal of Environmental Science and Technology*. 2009;3:042-049.
45. Obahiagbon FI, Oboh H, Uwumarongie-Ilori GE, Agho I. Selenium contents of some commonly consumed nigerian vegetables and physiological implications. *Chem. Soc. Nigeria*. 2011;36.
46. Henschler D, Kirschner U. On the absorption and toxicity of selenium sulfide. *Archiv fur Toxikologie*. 1969;24:341-344.
47. Morss SG, Olcott HS. Absence of effect of tocopherol on acute oral toxicity of sodium selenite in the rat. *Proceedings of the Society for Experimental Biology and Medicine*. 1967;124:483-485.
48. Cummins LM, Kimura ET. Safety evaluation of selenium sulfide antidandruff shampoos. *Toxicology and applied pharmacology*. 1971;20:89-96.
49. Okwara EC, Meludu SC, Okwara JE, Enwere OO, Diwe KC, Amah UK, et al. Selenium, zinc and magnesium status of HIV positive adults presenting at a university teaching hospital in Orlu-Eastern Nigeria. *Nigerian journal of medicine: Journal of the National Association of Resident Doctors of Nigeria*. 2012;21:165-168.
50. Nwagha UI, Ogbodo SO, Nwogu-Ikojo EE, Ibegbu DM, Ejezie FE, Nwagha TU, et al. Copper and selenium status of healthy pregnant women in Enugu, southeastern Nigeria. *Nigerian journal of clinical practice*. 2011;14:408-412.
51. Babalola OO, Anetor JI, Adeniji AA. Assessment of selenium status of healthy adults in south western nigeria. *Asset: An International Journal (Series A)*. 2003;3:111-120.
52. Arinola OG, Charles-Davies MA. Micronutrient levels in the plasma of Nigerian females with breast cancer. *African Journal of Biotechnology*. 2008;7.
53. Atiba AS, Oparinde DP, Jimoh AK, Babatunde OA, Adelekan A, Adeola Niran-Atiba T. Oxidative stress and serum selenium in HIV patients on different antiretroviral regimen. *Greener J. Med. Sci*. 2012;2:163-167.
54. Oyewole OS, Adesina AD, Oke OT, Tijani YO. Evaluation of essential trace elements in female type II diabetic patients in Nigerian population. *African Journal of Biotechnology*. 2014;13:1910-1914.
55. Duffield-Lillico AJ, Reid ME, Turnbull BW, Combs GF, Jr Slate EH, Fischbach LA, et al. Hypert thyroidism and Hypothyroidism. *American Association of Clinical Endocrinologists*. 2003;462-465.
56. Penney KL, Schumacher FR, Li H, Kraft P, Morris JS, Kurth T, et al. A large prospective study of SEP15 genetic variation, interaction with plasma selenium levels, and prostate cancer risk and survival. *Cancer Prevention Research*. 2010;3:604-610.
57. Durosinmi MA, Howard A, Akinola WO, Onayemi O. Selenium and aspirin in people living with HIV and AIDS in Nigeria. *The Nig Postgrad Med J*. 2008;15:215-217.
58. Famurewa AC, Akinosun OM. 'Plasma levels of α -tocopherol, -tocopherol and selenium in patients with prostate cancer in Nigeria'. *International Journal of Medicine and Biomedical Research*. 2014;3.
59. Moore JA, Noiva R, Wells IC. Selenium concentrations in plasma of patients with arteriographically defined coronary atherosclerosis. *Clinical Chemistry*. 1984;30:1171-1173.
60. Ajayi GO, Omilabu SA. Magnesium, selenium, zinc, copper and CD4 count in HIV1-and 2-positive Nigerians attending prepregnancy class at Prenatal Diagnosis and Therapy Centre of Lagos University. *Trace Elements and Electrolytes*. 2003;20:203-207.
61. Akinboro AO, Mejiuni DA, Onayemi O, Ayodele OE, Atiba AS, Bamimore GM. Serum selenium and skin diseases among Nigerians with human immunodeficiency virus/acquired immune deficiency syndrome. *HIV/AIDS (Auckland, NZ)*. 2013;5:215.

62. Luty-Frackiewicz A. The role of selenium in cancer and viral infection prevention. *International Journal of Occupational Medicine and Environmental Health*. 2005;18:305-311.
63. Eze JI, Okeke MC, Ngene AA, Omeje JN, Abonyi FO. Effects of dietary selenium supplementation on parasitemia, anemia and serum proteins of *Trypanosoma brucei brucei* infected rats. *Experimental Parasitology*. 2013;135:331-336.
64. Ukperoro JU, Offiah N, Idris T, Awogoke D. Antioxidant effect of zinc, selenium and their combination on the liver and kidney of alloxan-induced diabetes in rats. *Mediterranean Journal of Nutrition and Metabolism*. 2010;3:25-30.
65. Sadau Y, Adelaiye AB, Magaji RA, Ayo JO, Mabrouk MA, Isa AI. Role of Selenium and Vitamin E on Gastric Mucosal Damage Induced By Water-Immersion Restraint Stress in Wistar Rats. *IOSR J. Pharm. Biol. Sci*. 2015;10:34-39.