

The Occlusal Status of 6 to 12 Years Old Saudi Arabian Children: A Cross-sectional Study

Noura Abdulaziz Abumelha¹, Rawan Hamad Alyami², Nujud Shaya AlEdrees², Alanoud Saeed Abu Hatlah², Bayan Dleem Almoghamer² and Rafi Ahmad Togoo^{2*}

¹Department of Orthodontics, King Khalid University College of Dentistry, Abha, Saudi Arabia; ²Department of Pediatric dentistry & Orthodontics, King Khalid University College of Dentistry, Abha, Saudi Arabia

Corresponding author:

Prof. Rafi Ahmad Togoo,
Department of Pediatric Dentistry &
Orthodontics,
King Khalid University College of Dentistry,
Abha, Saudi Arabia,
Tel: +966542709996
E-mail: ratogo@kku.edu.sa

Abstract

Background: Malocclusion may cause frequent biting of the inner cheeks or tongue, aesthetic issues, discomfort when chewing or biting, speech problems, or development of mouth breathing therefore affecting the overall quality of life. Aim: To estimate the prevalence of different degrees of malocclusion associated dental complications among children aged 6-12 years. **Materials & Method:** A cross-sectional study was conducted among 526 children aged 6-12 years old after consents obtained from their parents. The Saudi children were selected from certain cities using proportional allocation technique by dividing these cities into districts where children were selected from malls and gardens with high attendance rate. **Results:** Based on molar relation, the prevalence of malocclusion in children aged 6-12 (children with mixed dentition) was 35.4%, of them 55% had class II division I with or without subdivisions and 34% had class III with or without subdivisions. The remaining 11% of children had different types of malocclusion include class II division II and combination of class II and class III malocclusions. Normal bite was reported in 34.8% of the children, other types of bites include open bite, deep bite, edge to edge bite which were recorded in 40.1%, 21.3%, 3.8% of children respectively. **Conclusion:** This study found more than one third of Saudi children affected by different types and degrees of malocclusion, malalignment and crowding. Significant associations were detected between age of children and prevalence of malocclusion and crowding.

Keywords: Children; Crowding; Diastema; Malocclusion; Prevalence; Saudi

Introduction

It has been demonstrated that type of occlusion of deciduous teeth is important in determining the occlusion in permanent teeth.^[1] The treatment of malocclusion has been a major challenge to dental professionals given the complicated and costly treatment involved, thus the emphasis is on early diagnosis and prevention. Determining prevalence of malocclusion in children leading to early diagnosis and establishing societal need for treatment helps orthodontists and pedodontists in devising a proper treatment plan and planning preventive and interceptive protocols.^[2,3]

Several studies have been published about occlusion of primary dentition in many countries,^[4-10] however in Arab countries the amount of literature that has been published is limited.^[11-13] Occlusion in primary dentition was found affected by various types of malocclusion, an old study noticed that as high as three quarters of 3-year-age children acquire flush occlusal relationship.^[4] Clinic reported a prevalence of 43% of asymmetric molar relationship among 61 children.^[14] Similarly, other Indian and Israeli studies reported more than half of children had flush terminal relationship with prevalence of 52% and 68% respectively.^[9,15] In Saudi children, a study found a prevalence of flush terminal relationship as high as 80% among 3-5 years old children.^[16] In addition, other malocclusions such as mesial and distal step have reported among Saudi children. Malocclusion refers to a number of possible conditions. It could be improper alignment of the teeth, or alteration in the appearance of the face. Malocclusion may cause frequent biting of the inner cheeks or tongue, aesthetic issues, discomfort when chewing or biting, speech problems, or development of mouth breathing therefore affecting the overall quality of life.^[17]

This study aimed to estimate the prevalence of different degrees of malocclusion associated dental complications among children aged 6-12 years. The findings of this study can help us to understand the societal need for treatment and helping all stakeholders to devise preventive measures and treatment plans accordingly.

Methods

A cross-sectional study was conducted among 526 children aged 6-12 years old after obtaining the consents from their parents. The Saudi children were selected from certain cities using proportional allocation technique by dividing these cities into districts where children were selected from malls and gardens with high attendance rate. The examination disposable kits with pen, pencils and rulers were used to assess the occlusion status of the study participants. Subjects were categorized as having Class I, Class II and Class III malocclusion using Angle's Classification of molar relationship.

Inclusion/Exclusion criteria

All children of age 6 to 12 years, willing to participate in the study were included. Children with cranio-facial anomalies, those who were medically compromised and who had undergone any orthodontic treatment were excluded from the study.

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 3.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as the author is credited and the new creations are licensed under the identical terms.

How to Cite this Article: Abumelha NA, et al. The Occlusal Status of 6 to 12 Years Old Saudi Arabian Children: A Cross-sectional Study. *Ann Med Health Sci Res.* 2018;8:401-403

The diagnosis was reported in data collection sheet along with the demographic data of subjects. Data thus collected was entered into Microsoft excel sheet and put to statistical analysis using SPSS version 19 (statistical package of social science). Prior to the conduct of the study, ethical clearance was obtained from college of dentistry.




Figure 1: Main types of malocclusion in Saudi 6-12 aged children based on molar relation.

Table 1: Socio-demographics of included children.

Demographics	Frequency	Percent
Sex	Male	116 (22.1)
	Female	410 (77.9)
	Khamis	290 (55.1)
Residence	Abha	164 (31.2)
	Others	72 (13.7)
Father education level	Less than Secondary	106 (20.2)
	Secondary	191 (36.3)
	More than Secondary	229 (43.5)
Mother education level	Less than Secondary	177 (33.7)
	Secondary	106 (20.2)
	More than Secondary	243 (46.2)

Table 2: Distribution of malocclusions among children aged 6-12 years old in Asser region.

Category	Frequency	Percent
Occlusion	Malocclusion	186 (35.4)
	Normal occlusion	340 (64.6)
	Class 1	342 (65.0)
	Class 2 division 1	68 (12.9)
	Class 2 division 1 with subdivision	35 (6.7)
	Class 2 division 2	9 (1.7)
	Class 2 division 2 with subdivision	8 (1.5)
Type of malocclusions based on molar relation	class 3	44 (8.4)
	Class 3 with subdivision	20 (3.8)
	Normal Bite	183 (34.8)
	Open Bite	211 (40.1)
	Deep Bite	112 (21.3)
	Edge to edge	20 (3.8)
	Normal bite	395 (75.1)
Type of malocclusions based on vertical relation	Anterior crossbite	53 (10.1)
	posterior crossbite (unilateral)	50 (9.5)
	posterior crossbite (bilateral)	16 (3.0)
	posterior crossbite (total)	7 (1.3)
	Over Jet (anterior)	5 (1.0)

Table 3: Prevalence of teeth spacing and crowding in 6-12-year-age children in Aseer region.

Category	Frequency	Percent
Teeth crowding	Absent	334 (63.5)
	Present (Upper)	20 (3.8)
	Present (Lower)	116 (22.1)
	Present (Upper and lower)	56 (10.6)
	Absent	312 (59.3)
Diastema	Present (Upper)	134 (25.5)
	Present (Lower)	5 (1.0)
	Present (Upper and lower)	9 (1.7)
Premature tooth loss	Physiological (Upper)	66 (12.5)
	No	436 (82.9)
	Yes (Deciduous)	22 (4.2)
	Yes (Permanent)	62 (11.8)
	Yes (Deciduous and permanent)	6 (1.1)

Results

A total sample of 526 children aged between 6-12 years old was included in this study, where 78% were females and 22% were males. More than half of children were from Khamis while about a third of children were selected from Abha city [Tables 1-3]. Based on molar relation, the prevalence of malocclusion in Saudi children aged 6-12 (children with mixed dentition) was 35% represented by 184 children, of them 55% had class II division I with or without subdivisions and 34% had class III with or without subdivisions. The remaining 11% of children with malocclusions had different types of malocclusion include class II division II and combination of class II and class III malocclusions [Figure 1].

The vertical relation between upper and lower dentations, normal bite was reported in 34.8% of the children, other types of bites include open bite, deep bite, edge to edge bite which were recorded in 40.1%, 21.3%, 3.8% of children respectively. The transverse relation between dentations was normal in 75.1% of children, while 10.1% and 13.8% of children had anterior and posterior cross bite respectively. The prevalence of diastema and tooth crowding were 40.7% and 36.5% respectively, in the subjects.

Only children aged 10-12 years old had significantly higher malocclusion prevalence than 6-9 years old children (p=0.011). Other associations between demographics of studied children and the presence of malocclusion were found statistically non-significant.

Discussion

Several studies have reported a high prevalence of malocclusion among school children which varied between populations. In Saudi Arabia, the seeking for orthodontic treatment has been grown in the last decade.^[18]

In the present study, the antero-posterior molar relationship showed the prevalence of 35% of malocclusion in Saudi children aged 6-12. Similarly, Alzubair and Ghandour found 30.6% prevalence of discrepancies in antero-posterior molar relationship in 12-year-age Yemeni children.^[12] Johannsdottir et al. found 33% and 36% prevalence of malocclusions in 6-year-age children.^[19] A study among Lithuanian 7-15 years aged children found that only 15.3% had normal occlusion.^[20] Thailander found that malocclusions found in 88% of 5-7 years old children. This difference in malocclusion prevalence can be attributed to the genetic factors and different criteria used to define malocclusions.

In the present study, 22.8% of studied children had class II and 12.2% had class III, while Johannsdottir et al. found 27% of boys and 32% of girls aged 6 years old had class II in Iceland, while 6.2% and 4.8 had class III in boys and girls respectively.^[19] Thilander et al. found a 21% and 3.7% prevalence of class II and class III malocclusion in 5-17 years old students^[21]. In Lithuanian children, class I, class II and class III occlusions were seen in 68.4%, 27.7% and 2.8% of them.^[20]

The present study found normal bite in 34.8% of the children, other types of bites include open bite, deep bite, edge to edge bite which were recorded in 40.1%, 21.3%, 3.8% of children respectively. Similar results of Thilander et al. who found 21.6% prevalence of deep bite in 5-17-years-old boys.^[21] These findings were different in Lithuanian children where the prevalence of deep bite (>3.0 mm) was 14.5% and the prevalence of open bite was only 3.5%.^[20] The transverse relation between dentitions in the studies children were normal in 75.1% of children, while 10.1% and 13.8% of children had anterior and posterior cross bite respectively. In Lithuania, posterior cross bite was found in 8.8% of schoolchildren.^[20]

In the present study, the prevalence of diastema was 40.7%, while Alzubair and Ghandour found 14.2% prevalence of diastema with 1 mm or more in 12 years children. Thilander et al. reported a prevalence of 13.5% in early mixed dentition that have been decreased with dentition development to be 3.7% in adolescents.^[21] Another study among Swedish school children found 5% prevalence of medial diastema. The prevalence of sucking habits among Saudi children was found as high as 48.36%,^[22] which explained to some extent the high prevalence of spacing in upper dentition. The prevalence of tooth crowding in our study was 36.5% while Alzubair and Ghandour found 31.4% prevalence of crowding.^[12] Thilander et al. found the crowding as the most common problem affected more than half of the studied 5-17 years aged students.^[21] In Lithuanian children, the prevalence of crowding in the upper and lower dentitions was 38.4% and 35.5% respectively. This prevalence was found significantly increased with age of children.^[20] This is in accordance with our study that found older children had significantly higher malocclusion prevalence than younger children.

Conclusion

This study found more than one third of Saudi children affected by different types and degrees of malocclusion, mal-alignment and crowding. Significant associations were detected between age of children and prevalence of malocclusion and crowding. These findings will highlight the high treatment needs of orthodontic treatment among Saudi children.

Acknowledgements

The authors would like to express gratitude to King Khalid University, Abha for its constant support in accomplishment of this study.

Conflict of Interest

The authors disclose that they have no conflicts of interest.

References

- Radnizic D. Dental crowding and its relationship to mesiodistal crown diameters and arch dimensions. *Am J Orthod Dentofacial Orthop* 1988;94:50-56.
- Al-Emran S, Wisth PJ, Boe OE. Prevalence of malocclusion and need for orthodontic treatment in Saudi Arabia. *Community Dent Oral Epidemiol* 1990;18:253-255.
- Almasri M. The prevalence of skeletal malocclusion in the Southern Aseer Region of Saudi Arabia. *American Journal of Medical Sciences and Medicine* 2014;2:13-15.
- Chapman H. The normal dental arch and its changes from birth to adult. *Br Dent J* 1935;58:201-229.
- Clinch LM. A longitudinal study of the mesiodistal crown diameters of the deciduous teeth and their permanent successors. *Eur Journal Orthod* 2007;29: i75-i81.
- Jensen E, Kai-Jen Yen P, Moorrees CF, Thomson SO. Mesiodistal crown diameters of the deciduous and permanent teeth in individuals. *J Dent Res* 1957 36:39-47.
- Boyko D. The incidence of primate spaces in fifty 3-year-old children of the Burlington study. *Am J Orthod Dentofacial Orthop* 1968;54:462-465.
- Otuyemi OD, Sote EO, Isiekwe MC, Jones SP. Occlusal relationships and spacing or crowding of teeth in the dentitions of 3–4-year-old Nigerian children. *Int J Paediatr Dent* 1997;7:155-160.
- Kaufman A, Koyoumdjisky E. Normal occlusal patterns in the deciduous dentition in preschool children in Israel. *J Dent Res* 1967;46:478-482.
- Margetts B, Brown T. Crown diameters of the deciduous teeth in Australian Aborigines. *Am J Phys Anthropol* 1978;48:493-502.
- Abu Alhajja E, Qudeimat M. Occlusion and tooth/arch dimensions in the primary dentition of preschool Jordanian children. *Int J Paediatr Dent* 2003;13:230-239.
- Al-Zubair NM, Ghandour IA. Occlusal status among Yemeni children. *APOS Trends in Orthodontics* 2014;4:93.
- Meer Z, Sadatullah S, Mohammad AW, Stephen AO, Pervez AR, Abdel BM. Prevalence of malocclusion and its common traits in Saudi males of Aseer region. *Journal of Dental Research and Review* 2016;3:99-102.
- Clinch LM. An analysis of serial models between three and eight years of age. *D Record* 1951;71:61.
- Nanda RS, Khan I, Anand R. Age changes in the occlusal pattern of deciduous dentition. *J Dent Res* 1973;2:221-224.
- Farsi NM, Salama FS. Characteristics of primary dentition occlusion in a group of Saudi children. *Int J Paediatr Dent* 1996;6:253-259.
- Bernabe E, Flores-Mir C, Sheiham A. Prevalence, intensity and extent of Oral Impacts on Daily Performances associated with self-perceived malocclusion in 11-12-year-old children. *BMC Oral Health* 2007;7:6.
- Hassan AH. Orthodontic treatment needs in the western region of Saudi Arabia: A research report. *Head & Face Medicine* 2006;2:2.
- Johannsdottir B, Wisth PJ, Magnusson TE. Prevalence of malocclusion in 6-year-old Icelandic children: a study using plaster models and orthopantomograms. *Acta Odontol Scand*, 1997;55:398-402.
- Sidlauskas A, Lopatiene K. The prevalence of malocclusion among 7–15-year-old Lithuanian schoolchildren. *Medicina* 2009;45:147-152.
- Thilander B, Pena L, Infante C, Parada SS, De Mayorga C. Prevalence of malocclusion and orthodontic treatment need in children and adolescents in Bogota, Colombia. An epidemiological study related to different stages of dental development. *Eur j orthod* 2001;23:153-167.
- Farsi NM, Salama FS. Sucking habits in Saudi children: Prevalence, contributing factors and effects on the primary dentition. *Pediatr Dent* 1997;19:28-33.