

Zinc Concentration during Pregnancy and Lactation in Enugu, South-East Nigeria

Ejezie FE* and Nwagha UI**

*Micronutrients/Molecular Toxicology Unit, Department of Medical Biochemistry, College of Medicine, University of Nigeria, Enugu Campus, Enugu State, Nigeria.

**Reproductive Biology/Endocrinology unit, Department of Physiology/Obstetrics and Gynaecology, College of Medicine, University of Nigeria, Enugu Campus, Enugu State, Nigeria.

Abstract

Background: There is no doubt that micronutrient deficiency is common in our environment. Replacements have thus been instituted without adequate information on specific and peculiar micronutrient needs of our people. Zinc is an essential micronutrient which is prone to maternal depletion during pregnancy and lactation. Unfortunately, studies in that regard in Nigeria has received diminutive interest.

Objectives: To determine zinc concentration during pregnancy and lactation.

Methods: This is a cohort study involving one hundred and twenty (120) pregnant women attending the antenatal clinic of one tertiary and two secondary health care institutions in Enugu. Thirty-five (35) apparently healthy, non pregnant women were used as control subjects. Atomic Absorption Spectrophotometer (AAS) was used to determine the zinc levels in the serum of the mothers during pregnancy, postpartum and in the breast milk.

Results: Results showed that mean serum zinc levels were significantly decreased ($P < 0.0001$) in pregnancy when compared with non-pregnant control subjects. The levels decreased as gestation progressed, with the lowest concentration of serum zinc obtained during the third trimester. It was also observed that serum zinc levels which decreased in pregnancy, increased non-significantly ($P = 0.12$) in mothers postpartum. In the breast milk, zinc concentration decreased significantly ($P < 0.0001$) as lactation progressed with the highest content evident in colostrum.

Conclusion; The observed significant decreases in the levels of zinc during pregnancy and in breast milk places the mothers and their neonates at risk and thus, necessitate maternal supplementation. Dietary interventions such as food diversification and biofortification are recommended to improve dietary zinc intakes in pregnant and lactating mothers, and infants in this region.

Key words: Zinc; pregnancy; lactation: Enugu; Nigeria.

Received on 12/09/2010; revised on 28/10/2010; accepted on 28/10/2010.

Ann Med Health Sci Res Jan 2011; 1(1) 69-76

Introduction

Several epidemiological and prospective

investigators and practitioners have accepted that micronutrients have a major function in many periods of a woman's life, particularly during pregnancy and lactation.¹ Understandably, nutrient needs increase during pregnancy and lactation to support foetal growth and milk production respectively.² Indeed, micronutrient deficiencies during pregnancy are associated with low birth

Correspondence:

Dr Fidelis Ejezie

Department of Biochemistry, College of Medicine, University of Nigeria, Enugu Camp'us, Enugu State, Nigeria.

E-mail: fidelis.ejezie@unn.edu.ng

weight, poor growth and development and sometimes neonatal and infant mortality. During lactation, a major reduction in the concentration of some of these nutrients in breast milk could occur and subsequently leads to infant depletion.³ Recent estimates indicate that globally, over two billion people are at risk for micronutrient deficiencies, despite recent efforts at prevention and management.⁴ These deficiencies are more pronounced during pregnancy and lactation, and in young children of Southeast Asia and sub-Saharan Africa.⁴

Zinc is an essential trace element in human and animal nutrition, with protective properties against endogenous or exogenous aggression, necessitated by the hyperproduction of highly reactive oxygen derivatives. It is an antioxidant, and also an effective 'scavenger', mopping up free radicals and reactive oxygen species.⁵ It is fundamental for growth, development, reproduction and immune response.⁶ Variations in plasma zinc concentration occasioned by disease, pregnancy, or stress and fluctuations in dietary intake are common.⁷ Zinc deficiency is estimated to be one of the commonest nutritional problems in the world.⁸ Indeed zinc deficiency in the female can lead to impaired synthesis/secretion of follicle stimulating and luteinising hormone, abnormal ovarian development, recurrent abortion, prolonged pregnancy and labour, teratogenicity, still births, pre-eclampsia, and low birth weight of infants.⁹

Human breast milk is a complex fluid, rich in nutrients and non-nutritional bioactive components.¹⁰ It is not only a source of nourishment but also a powerful antimicrobial agent.¹¹ The quality of a mother's breast milk is very crucial for the optimal survival of the new born baby. In our environment, maternal

malnutrition is common, and this has led to wild spread suggestions that routine micronutrient supplementation should be the rule. This supposition has led to the haphazard importation, into the Nigerian market, of numerous multivitamin and trace element compositions, with each drug company asserting superiority of its products over the others. Regrettably, there is paucity of information on maternal zinc status during pregnancy and lactation in our environment. Invariably, this has led to the production of supplements based on information that is completely extraterrestrial to our peculiar environment. In this regard, we decided to establish the baseline zinc status of our pregnant and lactating mothers to assist health care providers determine the actual amount of zinc required during pregnancy and lactation.

Subjects and Methods

Study Area

Created in 1991 from the old Anambra state, Enugu State is one of the 36 states of the Federal Republic of Nigeria. It has an approximate land mass of 8727.1km². It shares borders with Abia state to the south, Ebonyi state to the East, Benue state to the North East, Kogi state to the North West and Anambra state to the West. It has a mixed rural and urban population with majority being Igbo's, with a projected population of 3.3 million out of which about 50% are females. Enugu State has a crude birth rate of 45 per 1000, crude death rate of 18 per 1000 of the population and a life expectancy of 51years.¹² The maternal mortality rate ranges between 750 and 850 per 100,000 life births.^{13, 14} Enugu city is located in the hilly tropical environment about 230 m above sea level. The average annual temperature is between 23.1°C and 31°C with a rainfall of 1520 to 2030mm. There are two major seasons, rainy

season (April to October) and dry season (November to February). The city has a population of about 464,514 inhabitants.

Study Design and Setting

This is a cohort survey, undertaken between March and December 2009. After obtaining ethical clearance and informed verbal consent, 120 pregnant women, 40 in each trimester; first (0-13 weeks), second (14-27 weeks) and third (28 weeks and above) were randomly recruited. They were attending the ante-natal clinic of one tertiary and two secondary health care institutions in Enugu. The selection was based on lucky dip of "YES" or "NO". Women who had multiple pregnancies, obstetric haemorrhage, anaemia, fever, HIV positive, sickle cell disease, diabetes mellitus, malignancy, chronic alcoholics, tuberculosis, hypertension and any diagnosed inflammatory disorder in pregnancy were excluded. Also excluded were smokers and women on zinc supplements. The control group was made up of age and socioeconomic status matched 35 non-pregnant, non-lactating, apparently healthy mothers. They were recruited from the staff of UNTH and University of Nigeria, Enugu Campus (UNEC). The age range of the pregnant women was 19-42 years. Among those recruited, only 80 completed the study; twenty two (22) from the 1st trimester, 31 from the second trimester and 27 from the 3rd trimester.

Approval for the study was granted by the Ethics committee of the Hospital (ref: UNTH/CSA.329/Vol.6).

Methods

Socio-Economic Status of Subjects

Information on the socio-economic status was obtained by the administration of semi-structured, self-administered questionnaire. The socio-economic class was determined by the method of Szreter.¹⁵

Dietary Evaluation of Subjects

A 24-hour dietary recall interview was conducted for all the subjects to estimate their dietary zinc and calorie intake. In this method, the subjects were required to recall the individual's exact food intake during the previous twenty-four-hour period or preceding day. Detailed descriptions of all foods and beverages consumed, including cooking methods and brand names (where possible) were recorded. The amounts of nutrient intakes were calculated based on the nutrient composition of commonly eaten foods in Nigeria and other parts of the world.^{16,17}

Anthropometric/Socio-Demographic Characteristics.

The anthropometric/socio-demographic characteristics were obtained from the medical history, and physical examination. The weights were measured to the nearest 0.5 kilogram using a standard weighing scale (**STADIOMETER, SECA, MODEL 220, GERMANY**). The heights were measured in meters, with the same equipment without shoes, with the feet together, standing as tall as possible with the eyes level and looking straight ahead. Obstetric examinations, routine laboratory tests and ultra-sonography were performed.

Blood Sample Collection

About 5ml venous blood from the antecubital vein was collected from subjects at each period (pregnancy, and postpartum), using sterile, disposable syringes. The samples were transferred into sterile, anti-coagulant-free glass sample containers (plain tubes). Samples for the non pregnant group were collected on the 5th day of their menstrual cycle. The blood samples were allowed to stand for about thirty (30) minutes to clot and then centrifuged at 3,500 rpm for 15 minutes. The serum was collected and kept frozen at -20°C until analysed.

Breast Milk Sample Collection

Breast milk sample collection was carried out on the 3rd, 6th, 9th, 12th and 15th day after delivery since lactation did not commence on the day of delivery for most of the mothers. Breast milk samples (10ml) were collected from the lactating mothers by manual expression of the milk directly from the breast into sterile, plastic sample containers. After collecting the first 5ml, the baby was breastfed for 15 minutes and thereafter, an equal volume of hind-milk from the same breast was collected. After mixing thoroughly, the samples were immediately stored and frozen at -20°C prior to analysis. All milk samples were collected within the same period of the day (9.00 – 11.00 am) to avoid any possible diurnal changes.

Analysis of Zinc Concentration

Analyses (serum and milk) were done within two weeks of sample collection using Atomic Absorption Spectrophotometer (AAS), (Buck Scientific AAS/AES Model 205, United States of America).

Statistical Analysis

Analysis of data was done using SPSS version 11. Values were recorded as mean and standard deviation. D'Agostino & Pearson omnibus normality test was performed and data assumed Gaussian distribution. Consequently, a one way analysis of variance (ANOVA) followed by Scheffe' s post hoc tests. *P* values < 0.05 were considered significant.

Results

Table 1 represents the mean values of some anthropometric/sociodemographic characteristics of the subjects. The estimated daily zinc and caloric intake of the subjects did not vary significantly (*P* =0.54 and 0.62 respectively). The mean zinc level (µg/dl) in the control subjects was 109.75 ± 17.69. This decreased to 106.82 ± 19.76 in the 1st trimester, 101.65 ± 14.47 in the 2nd trimester and 85.57 ± 15.50 in the 3rd trimester. However immediately after delivery, the value rose to 90.66 ± 12.94. These values are statistically significant. (*P* <0.0001). However, after post hoc analysis, it was shown that the observed significant change is due to differences between controls versus 2nd trimester, control versus 3rd trimester and control versus postpartum. The level of zinc (µg/dl) in the breast milk (table 2) decreased significantly from 71.52 ± 4.06 on 0-3 days, 65.55 ± 3.74 4-6 days, 61.32 ± 3.14 7-9 days, 58.76 ± 2.73 10-12 days and 54.18 ± 3.44 13-15 days. (*P* <0.0001). Again, after post hoc analysis, it was observed that the differences between 7-9days and 10 -12days did not contribute to the statistically significant change.

Table 1: Some Anthropometric, Socio-Demographic Characteristics and Estimated Daily Nutrient Intake.

Variables	1 st trimester	2 nd trimester	3 rd trimester	Control
Mean age (years)	30.12±5.60	28.23±5.47	29.04±4.04	28.20±6.30
Mean gestational age (weeks)	9.94±1.75	21.27±3.83	34.42±2.90	0.000
Mean parity	1.00±1.41	1.23±1.74	1.85±1.46	1.53±1.77
Height(m)	1.65±.07	1.60±.08	1.66±.05	1.64±.05
Mean weight (kg)	69.71±14.10	71.14±14.77	81.46±10.35	64.60±10.62
EDCI (kcal/day)	2130±110	2200±110	2210±100	2100±108
EDZnI (mg/day)	26.65±7.10	24.92±6.70	23.43±5.20	21.72±4.90

EDCI (Estimated Daily Calorie Intake)

EDZnI (Estimated Daily Zinc Intake)

Table 2: Mean Breast Milk Zinc Concentrations (µg/dl) at Different Days of Postpartum/Lactation Period

Days Postpartum	Number of Subjects (n)	Mean ± SD
Day 0-3	10	71.52 ± 4.06
Day4 -6	14	65.55 ± 3.74
Day 7-9	19	61.32 ± 3.14
Day 10-12	22	58.76 ± 2.73
Day 13-15	15	54.18 ± 3.44

Values decreased as duration of lactation increased

Discussion

There is no doubt that zinc status is an imperative nutritional parameter during pregnancy and lactation.¹⁸ Accordingly, the determination of its levels has become a valuable tool in the management of pregnancy and its complications. The results of this study revealed that there was a statistically significant decrease in mean serum zinc concentration in pregnancy when compared with non-pregnant subjects. The decrease was progressive with the lowest mean serum zinc levels obtained in the third trimester. Significantly lower serum zinc levels in pregnancy have been reported in earlier studies in different parts of the world.^{18, 19, 20}

There is no doubt that the body's need for zinc is greatest during pregnancy, lactation, infancy and adolescence.⁹ Consequently, the need to increase zinc intake during pregnancy in this population through supplementation cannot be overemphasised. Several studies have continued to associate zinc deficiency with obstetric complications and hence continued to advocate increase in zinc intake during pregnancy and lactation.^{21, 22} Indeed it has been reported that oxidative stress can result from deficiency of trace elements such as zinc, copper and selenium.²³ Invariably, zinc supplementation reduces pregnancy complications, improves pregnancy

outcome, foetal development, lactation and neonatal growth.²¹

In this study, it was established that serum zinc concentration which decreased in pregnancy, increased slightly, though non-significantly in mothers after the baby was delivered. The slight increase could result from the expulsion of the placenta which transports maternal zinc to the foetus. Furthermore, it has been shown that zinc absorption efficiency increases markedly in early lactation compared with preconception levels.² Serum zinc homeostasis during pregnancy has been linked to placental hormones. As lactation commences, zinc is transferred from the mother to the breast milk.

In the present study, a progressive decline in the zinc concentrations of breast milk as the duration of lactation increased was demonstrated, with the highest concentration in colostrum (day 0-3 postpartum) and the least in mature milk (day 13-15 postpartum). The result is consistent with other reports which affirmed that concentrations of zinc in breast milk decreased significantly at successive stages of lactation.^{24, 25} Indeed it has been demonstrated that the macronutrient, mineral and trace element composition of breast milk fluctuates to a great extent as the duration of lactation increased.²⁴ Furthermore, breastfeeding routine/pattern; parity, age, and other maternal characteristics; regional differences, and in some situations, season of the year and maternal diet have been known to influence the concentration of breast milk constituents.²⁶

This study forms a hinge in the establishment of serum zinc reference levels in pregnancy, postpartum and breast milk of mothers in this region. As a result, it should constitute the

foundation for the formulation of sufficient micronutrient replacement policy for our pregnant and lactating mothers. Conversely, it would have been more appropriate if we had recruited all the women in the first trimester and followed them up until delivery. Also there may have been some bias in the assessment of zinc intake from the dietary recall. Moreover, one month dietary recall, wherever practicable gives superior dietary synopsis than a 24 hour recall. These will be addressed in further studies.

Our priority in developing countries should therefore be institution of a suitable strategy for the prevention and management of maternal undernutrition. This may require multiple interventions supported and promoted by different delivery strategies. Food supplements should be fortified and micronutrients formulated to suit our peculiar geographic and socioeconomic circumstances. This will go a long way to reduce the burden of maternal and infant morbidity and mortality. Furthermore, women should be educated on the importance of adequate and optimal macro and micronutrient intake during pregnancy and lactation.

References

1. Kontic-Vucinic O, Sulovic N and Radunovic N. Micronutrients in women's reproductive health: II. Minerals and trace elements. *Int J Ferti Women's Med* 2006; 51(3): 116-124.
2. King JC. Effect of reproduction on the bioavailability of calcium, zinc and selenium. *J Nutr* 2001; 131: 1355S-1358S.
3. Allen LH and Graham JM. Assuring micronutrient adequacy in the diets of young infants. In: Delange FM and West KPJ (Eds). *Micronutrient deficiencies in the first six*

- months of life. Basel: Vevey/S.Karger AG; 2003.
4. Ramakrishnan U. Prevalence of Micronutrient malnutrition worldwide. *Nutr Rev* 2002; 60: 46-52.
 5. Gutteridge JMC. Antioxidants, nutritional supplements and life-threatening diseases. *Br J Biomed Sci* 1994; 51: 288-295.
 6. Kornberg A. Origin of DNA on Earth. In: Supplement to DNA Replication. San Francisco: W.H. Freeman Co.; 1982; S224.
 7. Hambidge KM, Casey CE and Krebs NF. Zinc In: Mertz W. (Ed). Trace elements in human and animal nutrition, vol. 2, 5th ed. New York: Academic Press; 1986; 246.
 8. Prasad AS. Why a New Trace Element Journal? *J Trace Elem Exp Med* 1988; 1: 1-2.
 9. Bedwal RS and Bahuguna A. Zinc, copper and selenium in reproduction. *Cell Mol Life Sci* 1994; 50(7): 626-640.
 10. Prentice A. Breast milk calcium and phosphorus concentrations of mothers in rural Zaire. *Eur J Clin Nutr* 1991; 45:611-617.
 11. Bamgbose O, Opeolu BO and Bamgbose JT. Levels of Zinc in Breast Milk of Selected Nigerian Women in Abeokuta Township, Ogun-State. *Nig J Nutri Sci* 2008; 29(1): 153-160.
 12. State Ministry of Health, Enugu Health Sector Reform: Implementing the District Health System. Posted 2004. Assessed on 30th August 2010. Available from <http://www.enugustate.gov.ng/>.
 13. Onah HE, Okaro JM, Umeh U and Chigbu CO. Maternal mortality in health institutions with emergency obstetric care facilities in Enugu State, Nigeria. *J Obstet Gynecol.* 2005; 25 (6): 569-574.
 14. Ezugwu EC, Onah HE, Ezugwu FO and Okafor II. Maternal Mortality in a Transitional Hospital in Enugu, South East Nigeria. *Afr J Reprod Health* 2009; 13 (4):67-72.
 15. Szreter SRS. The Official representation of Social Classes in Britain, the United States and France: The Professional Model and "Les Cadres", Comparative Studies in Society and History 1993; 35, 2:285-317.
 16. Oguntano EB and Akinyele IO. Nutrient Composition of commonly eaten foods in Nigeria; raw, processed and prepared. Ibadan, Nigeria: Food Basket foundation Publication series; 1995.
 17. Paul AA and Southgate DAT (Eds). The composition of foods 4th ed. London: Her Majesty Stationery Office (HMSO); 1979.
 18. Izquierdo-Alvarez S, Castanon SG, Ruata ML, Aragues EF, Terraz PB, Irazabal YG, *et al.* Updating of normal levels of copper, zinc and selenium in serum of pregnant women. *J Trace Elem Med Biol* 2007; 21(Suppl 1): 49-52.
 19. Martin-Lagos F, Navarro-Alarcon M, Terres-Martos C, Lopez-Garcia de la Serrana H, Perez-Valero V and Lopez-Martinez MC. Zinc and Copper concentrations in serum from Spanish Women during pregnancy. *Biol Trace Elem Res* 1998; 61(1): 61-70.
 20. Arnaud J, Preziosi P, Mashako L, Galan P, Nsibu C, Favier A *et al.* Serum trace elements in Zairian mothers and their newborns. *Eur J Clin Nutr* 1994; 48(5): 341-348.
 21. Tamura T, Goldenberg RL, Johnston KE and DuBard M. Maternal plasma zinc concentrations and pregnancy outcome. *Am J Clin Nutr* 2000; 71(1): 109-113.

22. Ashraf M, Nasrullah M, Javeed S, Salam A, Ahmad Z and Rauf R. Maternal Serum Zinc level; its associate with complications of labor. *The Professional* 2001; 8(3): 383-386.
23. Al-Sayer H, Al-Bader A, Khoursheed M, Asfar S, Hussain T, Mathew A *et al.* Serum values of copper, zinc and selenium in adults resident in Kuwait. *Med Princ Prac* 2000; 9: 139-146.
24. Hannan MA, Dogadkin NN, Ashur IA and Markus WM. Copper, selenium, and zinc concentrations in human milk during the first three weeks of lactation. *Biol Trace Elem Res* 2005; 107(1): 11-20.
25. Yamawaki N, Yamada M, Kan-no T, Kojima T, Kaneko T and Yonekubo A. Macronutrient, mineral and trace element composition of breast milk from Japanese women. *Trace Elem Med Biol* 2005; 19(2-3): 171-181.
26. Domellöf M, Lönnerdal B, Dewey KG, Cohen RJ and Hernell O. Iron, zinc, and copper concentrations in breast milk are independent of maternal mineral status. *Am J Clin Nutr* 2004; 79:111–115.